

- Technology and equipment of food production

3/11 (81) 2016

Content

TECHNOLOGY AND EQUIPMENT OF FOOD PRODUCTION

- 4 Research of sedimentation stability of lipid-magnetite suspensions by the method of spectrophotometry
A. Alexandrov, I. Tsykhanovska, T. Gontar, N. Kokodiy
- 11 Developing a model of the foam emulsion system and confirming the role of the yield stress shear of interfacial adsorption layers to provide its formation and stability
A. Goralchuk, S. Omel'chenko, O. Kotlyar, O. Grinchenco, V. Mikhaylov
- 20 Optimization of formulation composition of the low-calorie emulsion fat systems
N. Tkachenko, P. Nekrasov, T. Makovska, L. Lanzhenko
- 27 Technology of specialty fats based on palm stearin
E. Kunitsa, O. Udovenko, E. Litvinenko, F. Gladkiy, I. Levchuk
- 33 The influence of mechanolysis on the activaton of nanocomplexes of heteropolysaccharides and proteins of plant biosystems in developing of nanotechnologies
R. Pavlyuk, V. Pogarska, T. Kotuyk, A. Pogarskiy, S. Loseva
- 40 Studying the accumulation of nitrogenous substances in biofortified pumpkin vegetables
G. Deinichenko, O. Yudicheva
- 46 Research into the impact of enzyme preparations on the processes of grain dough fermentation and bread quality
S. Oliinyk, O. Samokhvalova, A. Zaparenko, E. Shidakova-Kamenyuka, M. Chekanov
- 54 Consumer properties improvement of sugar cookies with fillings with non-traditional raw materials with high biological value
A. Tkachenko, I. Pakhomova
- 61 The effect of grape seed powder on the quality of butter biscuits
O. Samokhvalova, N. Grevtseva, T. Brykova, A. Grigorenko
- 67 Abstract&References

ГОЛОВНИЙ РЕДАКТОР

Бойник Анатолій Борисович

д. т. н., професор, Український державний університет залізничного транспорту (Україна)

Терзіян Ваган Якович

д. т. н., професор, Харківський національний університет радіоелектроніки (Україна)

професор Університету Ювяскюля (Фінляндія)

РЕДАКЦІЙНА КОЛЕГІЯ

ІНФОРМАЦІЙНІ ТЕХНОЛОГІЇ ТА СИСТЕМИ УПРАВЛІННЯ

Бойник А. Б., д. т. н., проф., Український державний університет залізничного транспорту, Харків (Україна); **Бутько Т. В.**, д. т. н., проф., Український державний університет залізничного транспорту, Харків (Україна); **Годлевський М. Д.**, д. т. н., проф., Національний технічний університет «Харківський політехнічний інститут», Харків (Україна); **Cardoso Jorge**, Professor of University of Coimbra, Faculty of Science and Technology, Coimbra (Portugal); **Omelayenko Borys**, PhD, Senior Software Engineer at Elsevier Amsterdam Area, Amsterdam (Netherlands); **Rab Nawaz Lodhi**, PhD, COMSATS Institute of Information Technology Sahiwal Campus (Pakistan); **Рибак Л. О.**, д. т. н., проф., Старооскольський технологічний інститут, Старий Оскол (Росія); **Самсонкін В. М.**, д. т. н., проф., Державний науково-дослідний центр залізничного транспорту України, Київ (Україна); **Соболєв Ю. В.**, д. т. н., проф., Український державний університет залізничного транспорту, Харків (Україна); **Terziyan Vagan**, Professor of University of Jyvaskyla, Department of Mathematical Information Technology, Jyvaskyla (Finland); **Фурман І. О.**, д. т. н., проф., Харківський державний технічний університет сільського господарства, Харків (Україна); **Jakab Frantisek**, Assoc. Professor of Technical University of Kosice, Department of Computers and Informatics, Kosice (Slovak Republic)

МАТЕМАТИКА І КІБЕРНЕТИКА - ПРИКЛАДНІ АСПЕКТИ

Ahmad Izhar, Associate Professor of King Fahd University of Petroleum and Minerals, Department of Mathematics and Statistics, Dhahran (Saudi Arabia); **Weber Gerhard Wilhelm**, Professor of Middle East Technical University, Institute of Applied Mathematics, Ankara (Turkey); **Дьомін Д. О.**, д. т. н., проф., Національний технічний університет «Харківський політехнічний інститут», директор ПП «Технологічний Центр», Харків (Україна); **Зелик Я. І.**, д. т. н., провідний науковий співробітник, Інститут космічних досліджень Національної академії наук України та Державне космічне агентство України, Київ (Україна); **Тєвяшев А. Д.**, д. т. н., проф., Харківський Національний університет радіоелектроніки, Харків (Україна); **Trujillo Juan J.**, Professor of Universidad de la Laguna, Faculty of Mathematics, San Cristobal de La Laguna (Spain)

ПРИКЛАДНА МЕХАНІКА

Andrianov Igor, Professor of RWTH Aachen University, Department of General Mechanics, Aachen (Germany); **Дудніков А. А.**, к. т. н., професор, Полтавська державна аграрна академія, Полтава (Україна); **Львов Г. І.**, д. т. н., проф., Національний технічний університет «Харківський політехнічний інститут», Харків (Україна); **Machado Jose Antonio Tenreiro**, Professor of Polytechnic of Porto, Institute of Engineering, Department of Electrical Engineering, (Portugal); **Пермяков О. А.**, д. т. н., проф., Національний технічний університет «Харківський політехнічний інститут», Харків (Україна); **Подригало М. А.**, д. т. н., проф., Харківський Національний автодорожній технічний університет, Харків (Україна); **Самородов В. Б.**, д. т. н., проф., Національний технічний університет «Харківський політехнічний інститут», Харків (Україна)

ПРИКЛАДНА ФІЗИКА

Гламаздін О. В., к. ф.-м. н., старший науковий співробітник, Національний науковий центр «Харківський фізико-технічний інститут», Харків (Україна); **Мар'янчук П. Д.**, д. ф.-м. н., проф., Чернівецький національний університет ім. Ю. Федьковича, Чернівці (Україна); **Новосядлій С. П.**, д. т. н., проф., Прикарпатський національний університет ім. Василя Стефаника, Івано-Франківськ (Україна); **Соболь О. В.**, д. ф.-м. н., проф., Національний технічний університет «Харківський політехнічний інститут», Харків (Україна); **Старіков В. В.**, к. ф.-м. н., старший науковий співробітник, Національний технічний університет «Харківський політехнічний інститут», Харків (Україна)

МАТЕРІАЛОЗНАВСТВО, ТЕХНОЛОГІЇ ОРГАНІЧНИХ І НЕОРГАНІЧНИХ РЕЧОВИН

Вахула Я. І., д. т. н., проф., Національний університет «Львівська політехніка», Львів (Україна); **Arvaidas Galdikas**, Professor Kaunas University of Technology, Department of Physics, Kaunas (Lithuania); **Глікін М. А.**, д. т. н., проф., Технологічний інститут Східноукраїнського національного університету ім. В. Даля, Сєверодонецьк (Україна); **Капустін О. Є.**, д. х. н., проф., Приазовський державний технічний університет, Маріуполь (Україна); **Carda Juan B.**, Professor of Universidad Jaume I, Department of Inorganic Chemistry, Castellon de la Plana (Spain); **Кондратов С. О.**, д. х. н., проф., Інститут хімічних технологій Східноукраїнського національного університету ім. В. Даля, Луганськ (Україна); **Михайлів В. М.**, д. т. н., проф., Харківський державний університет харчування і торгівлі, Харків (Україна); **Черевко О. І.**, д. т. н., проф., Харківський державний університет харчування і торгівлі, Харків (Україна); **Чумак В. Л.**, д. х. н., проф., Національний авіаційний університет, Київ (Україна)

ЕНЕРГОЗБЕРІГАЮЧІ ТЕХНОЛОГІЇ ТА ОБЛАДНАННЯ

Данько В. Г., д. т. н., проф., Національний технічний університет «Харківський політехнічний інститут», Харків (Україна); **Кліменко Б. В.**, д. т. н., проф., Національний технічний університет «Харківський політехнічний інститут», Харків (Україна); **Sutikno Tole**, Professor of Universitas Ahmad Dahlan, Department of Electrical Engineering, Yogyakarta (Indonesia); **Терещенко Т. О.**, д. т. н., проф., Національний технічний університет України «Київський політехнічний інститут», Київ (Україна)

EDITOR IN CHIEF

Boynik Anatoly

PhD, Professor of Ukrainian State University of Railway Transport (Ukraine)

Terziyan Vagan

PhD, Professor of Kharkov National University of Radioelectronics (Ukraine)

Professor of the University of Jyvaskyla (Finland)

EDITORIAL BOARD

COMPUTER SCIENCE

Boynik Anatoly, Professor of Ukrainian State University of Railway Transport, Department of Automation and Computer telecontrol traffic, Kharkov (Ukraine); **Butko Tatiana**, Professor of Ukrainian State University of Railway Transport, Department of operational work and international transportation, Kharkov (Ukraine); **Cardoso Jorge**, Professor of University of Coimbra, Faculty of Science and Technology, Coimbra (Portugal); **Furman Illya**, Professor of Petro Vasylchenko National Technical University of Agriculture, Department of Agricultural Mechanization, Kharkov (Ukraine); **Hodlyevskiy Mykhailo**, Professor of National Technical University «Kharkiv Polytechnic Institute», Department of Automated Control Systems, Kharkov (Ukraine); **Jakab Frantisek**, Assoc. professor of Technical University of Kosice, Department of Computers and Informatics, Kosice (Slovak Republic); **Omelayenko Borys**, PhD, Senior Software Engineer at Elsevier Amsterdam Area, Amsterdam (Netherlands); **Rab Nawaz Lodhi**, PhD, COMSATS Institute of Information Technology Sahiwal Campus (Pakistan); **Rybalk Larisa**, Professor of Staroskol Institute of Technology, Department of Automation and Industrial Electronics, Stary Oskol (Russia); **Samsomkin Valery**, Professor, Director of the State Research Center Railway Transport of Ukraine, Kyiv (Ukraine); **Sobolev Yuryi**, Professor, Advisor to the Rector of Ukrainian State University of Railway Transport, Kharkov (Ukraine); **Terziyan Vagan**, Professor of University of Jyvaskyla, Department of Mathematical Information Technology, Jyvaskyla (Finland)

MATHEMATICS

Ahmad Izhar, Associate Professor of King Fahd University of Petroleum and Minerals, Department of Mathematics and Statistics, Dhahran (Saudi Arabia); **Demin Dmitriy**, Professor of National Technical University «Kharkiv Polytechnic Institute», director of the Private Company «Technology Center», Kharkov (Ukraine); **Teviashev Andrew**, Professor of Kharkov National University of Radioelectronics, Department of Applied Mathematics, Kharkov (Ukraine); **Trujillo Juan J.**, Professor of Universidad de la Laguna, Faculty of Mathematics, San Cristobal de La Laguna (Spain); **Weber Gerhard Wilhelm**, Professor of Middle East Technical University, Institute of Applied Mathematics, Ankara (Turkey); **Zelyuk Yarema**, Leading Researcher of Space Research Institute of the National Academy of Sciences of Ukraine and National Space Agency of Ukraine, Kyiv (Ukraine)

ENGINEERING

Andrianov Igor, Professor of RWTH Aachen University, Department of General Mechanics, Aachen, (Germany); **Dudnikov Anatoly**, Professor of Poltava State Agrarian Academy, Department of the Repair machines and technology of constructional materials, Poltava (Ukraine); **Lvov Hennadiy**, Professor of National Technical University «Kharkiv Polytechnic Institute», Department of Dynamics and Strength of Machines, Kharkov (Ukraine); **Machado José António Teneiro**, Professor of Polytechnic of Porto, Institute of Engineering, Department of Electrical Engineering, (Portugal); **Permiakov Alexander**, Professor of National Technical University «Kharkiv Polytechnic Institute», Department of Mechanical Engineering, Kharkov (Ukraine); **Podrigalo Mykhailo**, Professor of Kharkiv National Automobile and Highway University, Department of Mechanical Engineering Technologies and Repairs, Kharkov (Ukraine); **Samorodov Vadim**, Professor of National Technical University «Kharkiv Polytechnic Institute», Department of Cars and Tractors, Kharkov (Ukraine)

APPLIED PHYSICS

Glamazdin Alexander, PhD, National Science Center «Kharkov Institute of Physics and Technology», Kharkov (Ukraine); **Novosiadliy Stepan**, Professor of Vasyl Stefanyk Precarpathian National University, Department of Physics and Technology, (Ukraine); **Maryanchuk Pavlo**, Professor of Yuriy Fedkovych Chernivtsi National University, Department of Physics, (Ukraine); **Sobol Oleg**, Professor of National Technical University «Kharkiv Polytechnic Institute», Department of Materials Science, Kharkov (Ukraine); **Starikov Vadim**, Senior Researcher of National Technical University «Kharkiv Polytechnic Institute», Department of Physics of metals and semiconductors, Kharkov (Ukraine)

MATERIALS SCIENCE, CHEMISTRY AND CHEMICAL ENGINEERING

Carda Juan B., Professor of Universidad Jaume I, Department of Inorganic Chemistry, Castellon de la Plana (Spain); **Cherevko Alexander**, Professor of Kharkiv State University of Food Technology and Trade, Kharkov (Ukraine); **Chumak Vitaliy**, Professor of National Aviation University, Department of Chemistry and Chemical Engineering, Kyiv (Ukraine); **Galdikas Arvaidas**, Professor of Kaunas University of Technology, Department of Physics, Kaunas (Lithuania); **Glikin Marat**, Professor of East-Ukrainian National University, Technological Institute, Department of Technology of organic substances, fuels, and polymers, Severodonetsk (Ukraine); **Kapustin Alexey**, Professor of Pryazovskyi State Technical University, Department of Chemistry, Mariupol (Ukraine); **Kondratov Sergey**, Professor of Volodymyr Dahl East-Ukrainian National University, Department of Mathematics and Computer Science, Lugansk (Ukraine); **Mihaylov Valerii**, Professor of Kharkiv State University of Food Technology and Trade, Kharkov (Ukraine); **Vakhula Yaroslav**, Professor of Lviv Polytechnic National University, Department of Silicate Engineering, Lviv (Ukraine)

ENERGY

Danko Vladimir, Professor of National Technical University «Kharkiv Polytechnic Institute», Department of Electrical Engineering, Kharkov (Ukraine); **Klimenko Boris**, Professor of National Technical University «Kharkiv Polytechnic Institute», Department of Electrical Apparatus, Kharkov (Ukraine); **Sutikno Tole**, Professor of Universitas Ahmad Dahlan, Department of Electrical Engineering, Yogyakarta (Indonesia); **Tereshchenko Tatiana**, Professor of National Technical University of Ukraine «Kyiv Polytechnic Institute», Department of Industrial Electronics, Kyiv (Ukraine)

<p>Honorary editor I. G. Filippenko Doctor of Technical Sciences, Professor Ukrainian State University of Railway Transport (Ukraine)</p> <p>Establishers PC «TECHNOLOGY CENTER» Ukrainian State University of Railway Transport</p> <p>Publisher PC «TECHNOLOGY CENTER»</p> <p>Editorial office's and publisher's address: Shatilova dacha str., 4, Kharkiv, Ukraine, 61145</p> <p>Contact information Tel.: +38 (057) 750-89-90 E-mail: eejet.kh@gmail.com Website: http://www.jet.com.ua, http://journals.uran.ua/eejet</p>	<p>Journal Indexing</p> <ul style="list-style-type: none">■ Scopus■ CrossRef■ American Chemical Society■ EBSCO■ Index Copernicus■ Российский индекс научного цитирования (РИНЦ)■ Ulrich's Periodicals Directory■ DRIVER■ Bielefeld Academic Search Engine (BASE)■ WorldCat■ Electronic Journals Library■ DOAJ■ ResearchBib■ Polska Bibliografia Naukowa■ Directory of Research Journals Indexing■ Directory Indexing of International Research Journals■ Open Academic Journals Index■ Sherpa/Romeo	<p>Свідоцтво про державну реєстрації журналу KB № 21546-11446 ПР від 08.09.2015</p> <p>Атестовано Вищою Атестаційною Комісією України Перелік № 12 постанови Президії ВАК № 1-05.36 від 11.06.03</p> <p>Постановою Президії ВАК України № 1-05/2 від 27.05.2009, № 1-05/3 від 08.07.2009 Бюлєтень ВАК України</p> <p>Наказом Міністерства освіти і науки України №793 від 04.07.2014</p> <p>Рекомендовано Вченю Радю протокол № 5 від 31.05.16 р.</p> <p>Підписано до друку 02.06.2016 р.</p> <p>Формат 60 × 84 1/8. Ум.-друк. арк. 9,5. Обл.-вид. арк. 8,84 Тираж 1000 екз.</p>
--	---	--

Наведені результати товарознавчих досліджень нових зразків печива цукрового з начинками «Дачне» та «Ясне сонечко». Нетрадиційна сировина, використана у рецепттурах печива, відрізняється збалансованими мінеральним та вітамінним складом, а також ефективною антиокислювальною здатністю, що дозволяє подовжити терміни зберігання продукції. Розроблені зразки печива цукрового відповідають нормативній документації за фізико-хімічними показниками, а також за показниками мікробіологічної та токсикологічної безпеки

Ключові слова: борошняні кондитерські вироби, нетрадиційна сировина, споживні властивості, харчова цінність, цукрове печиво з начинками

Приведены результаты товароведных исследований новых образцов печенья сахарного с начинками «Дачное» и «Ясное солнышко». Нетрадиционное сырье, использованное в рецептурах печенья, отличается сбалансированным минеральным витаминным составом, а также эффективным антиоксидантным действием, что позволяет продлить сроки хранения продукции. Разработанные образцы печенья сахарного соответствуют нормативной документации по физико-химическим показателям, а также по показателям микробиологической и токсикологической безопасности

Ключевые слова: муочные кондитерские изделия, нетрадиционное сырье, потребительские свойства, пищевая ценность, сахарное печенье с начинками

UDC 664.68
DOI: 10.15587/1729-4061.2016.70950

CONSUMER PROPERTIES IMPROVEMENT OF SUGAR COOKIES WITH FILLINGS WITH NON-TRADITIONAL RAW MATERIALS WITH HIGH BIOLOGICAL VALUE

A. Tkachenko

PhD, Assistant

Department of an expertise of goods and custom*

E-mail: alina_biaf@ukr.net

I. Pakhomova

PhD, Assistant

Department of commodity nonfood products*

E-mail: inpakhomova@gmail.com

*Poltava University of Economics and Trade

Kovalia str., 3, Poltava, Ukraine, 36014

1. Introduction

The pastry products, especially sugar cookies, are in steady demand in the adult and child population according to the marketing research [1]. But the traditional raw materials for preparation of sugar cookies are not full in the biological aspect because of high calorie, high content of saturated fat acids and low nutrition value. The improvement of consumer properties of sugar cookies with fillings is, therefore, acute. The usage of local resources of non-traditional raw materials is most promising and effective in the technological and economical aspects. The unconventional raw materials are a valuable source of healthy nutrients [2]. Therefore, an important task is the rational combination of different sorts of natural raw materials for optimization of the composition and improvement of consumer properties of sugar cookies with fillings [3].

The relevance of researches in this area is that by improving the nutritional value of sugar cookies it is possible to influence on the overall diet of the population because sugar cookies are in high demand according to the researches [1]. The population's modern nutrition imbalance encourages the creation of new food products with improved consumer properties. The considerable problem which needs to be solved is the improvement of amino acid, fatty acid, vitamin and mineral composition in the pastry products.

2. Analysis of previous studies and statement of the problem

Fat, wheat flour, eggs and sugar are traditional raw materials for the sugar cookies production. Analysis of the chemical composition of these ingredients, especially wheat flour [5] and fat [6] proves the need of the usage of non-traditional raw materials for the sugar cookies production. The developments of new rational recipes of pastry products with increased nutritional value with partial or full substitution of traditional ingredients on the milk-containing products, vegetable oils, processed fruit and vegetables and medical and industrial raw materials are conducted [7].

The improvement of the amino acid composition of pastry products can be achieved by the usage of milk products, processed legumes and oilseeds and protein enriching agents in their recipes. The scientists conduct developments for enriching the protein composition of sunflowers [8] and biological fortification of peanuts for optimization of their amino acid composition for confectionery production with increased protein's biological value. In particular, the lactic cheese and sweet whey powder are promising raw materials for the enriching pastry products with protein and minerals. But the question remains little known about the rational correlation of milk products in the recipes of pastry products. The scientific researches show improvement of amino acid composition of pastry products owing to the usage of

bean sprouts, pea grits [11], chickpeas flour and lentil flour in the recipes [12]. The replacement of 30 % wheat flour with the okara (mass produced by pressing bean milk on the filter-press) allows to enrich the product with amino acids [13]. As protein preparations for enriching the pastry products, egg whites enriched with proteins can be used. As a result, not only the amino acid composition is improved, the product acquires dietary properties [14].

The issue of the day in the food production is also the balancing of fatty acid composition of products. It is found that the fraction of unsaturated fatty acids increases and the fraction of saturated fatty acids decreases owing to the addition of non-traditional oils to recipes [15].

It is proved that non-traditional oils surpass the margarine in the content of polyunsaturated fatty acids [5]. The linseed oil has a balanced chemical composition and a high amount of polyunsaturated fatty acids. The amount of polyunsaturated fatty acids in the linseed oil is 5,38 times higher than in the margarine. The amount of polyunsaturated fatty acids in the soybean, pumpkin, sea buckthorn oils is 4,69; 4,61; 4 times higher than in the margarine [16]. The search for vegetable oils with high content α -linolenic acid which is useful for the organism is being conducted. These include Sacha Inchi oil, peony oil, sea buckthorn oil, cypress oil, cress salad oil. The content of α -linolenic acid is 50; 45; 35; 30 % in these oils [17]. The sea buckthorn oil effect on the fatty acid content of lipids is little known. However, it has been scientifically proven that sea buckthorn oil has a balanced ratio of omega-3 and omega-6 fatty acids. The sea buckthorn oil is the next after linseed oil by the content of α -linolenic acid. It allows considering oil as a valuable enricher for fat-containing products [18].

The considerable direction is enriching of confectionery with a limited content of macro- and micro elements and vitamins [19]. This need is dictated by objective environmental factors, which are associated with the change of the composition and nutritional value of products which are used. The next reason is the transformation of the lifestyle related to decreased physical energy expenditure [20].

A special attention from the scientific point of view must be paid to the enriching of fillings of pastry products because some vitamins are thermolabile. It is offered to include processed products of sugar beet [21], sea buckthorn concentrate [22], fruits chokeberry [23], etc. in the composition of fillings. The promising direction in the pastry production is the usage of medical and industrial raw materials and its processed products such as viburnum, hawthorn, buckthorn [24], goji berries [25]. The usage of these ingredients allows to enrich products with vitamins PP, B₁, B₂, mineral elements and cellulose.

Alternative kinds of flour, especially soybean and other legume crops have a positive impact on the improvement of nutritional value, particularly vitamin and mineral composition [26]. The soybean protein isolate and the meal of milk thistle in the recipes of custard cakes and cookies improve considerably the protein and mineral composition [27].

The developments using artichoke's powders [28], moringa and yam powders are proposed for the improvement of consumer properties of pastry products [29].

The scientific developments concerning the improvement of consumer properties of pastry products are varied, but the multifactorial impact of non-traditional oils and fruit and vegetable raw materials on the formation of nutritional

and biological value of food products is little studied [2]. Therefore, search for new recipes with non-traditional raw materials for the improvement of protein, fatty acid, vitamin and mineral composition of sugar cookies is relevant. The research presented in the paper aimed at the improvement of consumer properties and nutritional value of products for making better the diet of the population.

3. The purpose and objectives of the study

The purpose of the research is scientific and practical reasoning and developing the recipes of sugar cookies with fillings with the usage of the unconventional supplements to improve its consumer properties.

The purpose can be achieved by solving the following objectives:

- developing of the new recipes of sugar cookies;
- determination of the quality according to the standard;
- study of the amino acid, fatty acid, mineral and vitamin composition of sugar cookies;
- the comprehensive quality assessment.

4. Materials and Methods

The object of the study are the samples of the cookies "Dachne" and "Yasne sonechko" with the usage of the non-traditional raw materials.

The cookies "Dachne" has been produced with the usage of non-traditional raw materials: dried apples powders, dried raspberry leaf powders, dried calendula officinalis powders, pumpkin oil. The cherry plum and zucchini jam enriched with the preparation of the eggshell with lemon juice had been used for the production of fillings.

The cookies "Yasne sonechko" contains lungwort officinalis powders, dried apricot powders, sea buckthorn oil. The sea buckthorn jam with calendula syrup had been used for the production of fillings.

As a check sample, the cookies "Litne" have been used. It is the classic example of sugar cookies according to its recipe. For developing the recipes of sugar cookies, the mathematical modeling of prescription formulations has been used. Restrictions on the total content of ingredients in the recipe are determined by the formula:

$$\sum_{i=1}^j x_i = 1000, \quad (1)$$

where x_i , $i=1, 2, \dots, j$ is the unknown amount of raw materials of the i type (grams).

Process conditions of the required moisture content of the products are as follows:

$$0,05 \sum_{i=1}^j x_i \leq \sum_{i=1}^j \lambda_i x_i \leq 0,1 \sum_{i=1}^j x_i, \quad (2)$$

where x_i , $i=1, 2, \dots, j$ is the unknown amount of raw materials of the i type (grams); λ_i is the water content in 1 gram of the i ingredient.

Additional conditions of the projected product enrichment with nutrients were determined as a percentage relative to the daily needs.

The following objective function is advisable

$$Z = \sum_{i=1}^j a_i x_i \rightarrow \max, \quad (3)$$

where x_i , $i=1, 2, \dots, j$ is the unknown amount of raw materials of the i type (grams); a_n is the content of the nutrient of the n type in the 1 gram of the i ingredient (grams).

The task of optimizing the content of the ingredients in the new sugar cookies is the determination of the vector $X=(x_1, x_2, \dots, x_j)$, maximizing the objective function provided that the coordinates of the vector satisfy the system of inequalities and equations. The solution of the problem is obtained by the simplex method in the MathCAD (Prime 3.0) [31].

The sensory, physical and chemical and microbiological analysis methods were used for the study of consumer properties of new sugar cookies and change of its quality in the goods turnover process [32].

The amino acid composition of sugar cookies was determined by ion exchange liquid-column chromatography on the amino acids automatic analyzer T 339 ("Microtechnick, Czech Republic) [33].

The amino acid score was determined by the ratio of the relevant essential amino acid in the 1 gram of protein of the cookies to the required content in the "ideal protein" according to the FAO/WHO scale [34].

The fatty acid composition of sugar cookies was determined by gas chromatography method on the gas chromatograph HP 6890 (Agilent, CIIIA) [35].

The mineral composition of sugar cookies was determined by atomic absorption spectrophotometry method on the atomic absorption spectrophotometer C-115 PK (Semi, Ukraine) [35].

Retinol (vitamin A) was determined by the colorimetric method, which is based on the reaction retinol with antimony trichloride in chloroform with the formation of blue color. The intensity of the color is directly proportional to the content of vitamin A. Ascorbic acid (vitamin C) has been determined by the Tillmans method. Tocopherol (vitamin E) has been determined by liquid chromatography with high isolation capacity [35].

The conventional methods have been used for the study of the toxic elements content in the new cookies. Copper, zinc, lead and cadmium have been determined by the atomic absorption method; arsenic was determined by the colorimetric method, mercury was determined by the flameless atomic absorption method [35].

5. The results of the study of consumer properties of new sugar cookies with fillings

The selection of the formulation was carried out by mathematical modeling method and on the base of product's organoleptic properties. New ingredients have been selected according to their rational relation. Adjustment of the formulations has reduced the content of wheat flour and sugar in the cookies. The content of traditional and non-traditional raw materials is presented in Table 1. It should be noted that the raw materials which have been used for the production of the new cookies are produced in Ukraine and certified in accordance with current legislation.

According to the results of the tasting assessment of sugar cookies, which was held by the tasting commission consisting of 10 people at the Poltava University of Economics

and Trade (Ukraine), it has been found that the developed samples of sugar cookies with fillings are characterized by better organoleptic properties than the check sample. The organoleptic assessment was held using the 50-point scale, developed by the authors.

Table 1

The content of components in the new samples of cookies, kg/t

Ingredient	«Dachne»	«Yasne sonechko»
Wheat flour	526,03	608,07
Sugar powder	185,05	179,98
Invert syrup	28,02	29,71
Vegetable-cream mixture	102,10	106,67
Skim milk powder	—	36,05
Whey protein concentrate	52,20	63,16
Fat-free whey powder	21,36	—
Melange	25,69	23,93
Vanilla essence	3,15	2,00
Baking soda	4,14	4,80
Ammonium carbon salt	0,63	0,80
Salt	4,14	4,18
Dried raspberry leaf powder	11,72	—
Lungwort officinalis powder	—	2,60
Dried apple powder	63,31	—
Dried apricot powder	—	49,52
Dried calendula officinalis powder	2,86	—
Pumpkin oil	15,14	—
Sea buckthorn oil	—	22,40
Cherry plum and zucchini jam	124,45	—
Preparation of eggshell with lemon juice	9,43	—
Sea buckthorn jam	—	114,45
Calendula syrup	—	13,08

The taste and smell are the most important parameters for consumers [36]. According to these parameters, sugar cookies "Yasne sonechko" have got the most points (4,89 points). Sugar cookies "Dachne" have received the most points according to the parameter of "severity of additives". Sugar cookies "Yasne sonechko" have had attractive color. They have received by this parameter maximal 5 points. They have received by the appearance maximal 4,89 points.




Fig.1. Profilogramm of organoleptic properties of sugar cookies

The results of the study of physical and chemical indicators of the samples of sugar cookies (Table 2) show

that they comply with European regulations, especially Regulation of the European Parliament and of the Council «Laying down specific rules for the organisation of official controls on products of animal origin intended for human consumption».

Table 2
Physical-chemical indicators of new sugar cookies

Indicator	Norm	Research samples of cookies		
		Check sample	«Dachne»	«Yasne sonechko»
Humidity, %	10±2	9,00±0,29	9,50±0,27	6,10±0,29
Mass fraction of total sugars in terms of dry matter (by sucrose), %	not>27,0	25,16±0,05	21,55±0,05	22,23±0,05
Fat content in terms of dry matter, %	10±2,0	10,84±0,05	10,19±0,05	10,54±0,05
Alkalinity, degree.	not>2,0	1,6±0,05	1,4±0,05	1,2±0,04
Mass fraction of insoluble ash in solution with mass fraction of 10 % hydrochloric acid, %	not>0,1	0,06±0,01	0,02±0,01	0,02±0,01
Water absorption, %	not<150	165±7,90	177±9,90	177±9,90

According to the study, the bacteria *Escherichia coli*, pathogens, molds and yeasts have not been found in the fresh samples. The amount of mesophilic aerobic and facultative anaerobic microorganisms has not exceeded the norm. The amount of toxic elements has been in full record with the norm.

The nutritional and energy value has improved owing to adjustment of the recipes (Table 3).

Table 3
Nutritional and energy value of the new cookies

Samples of cookies	Content, g/100 g				Energy value kkal/100 g
	fats	proteins	carbohydrates	humidity	
Check sample	10,84	9,39	74,95	5,00	434,92
«Dachne»	10,19	9,41	70,00	9,50	409,35
«Yasne sonechko»	10,54	10,50	71,48	6,10	422,78

Table 3 shows significant increase of the protein content in the cookies «Yasne sonechko» (by 1,11 g/100 g) as compared with the check sample. The decrease of the carbohydrates content in all samples owing to the replacement of the main ingredients (wheat flour, sugar powder, invert syrup) with the other components is achieved. Especially it is shown in the cookies «Dachne» (by 4,95 g/100 g) as compared with the check sample. The cookies «Dachne» are characterized by a high amount of amino acids. It contains the whey protein concentrate. The total amount of amino acids is increased by 19,76 % compared to the check sam-

ple. The total amount of essential amino acids is increased by 34,83 % as compared with the check sample. The total amount of essential amino acids in the cookies «Yasne sonechko» is increased by 32,54 % as compared with the check sample.

One of the most valuable amino acids is lysine. Lysine deficiency in the diet leads to hemodyscrasia, reducing the number of red blood cells and reduction of hemoglobin in blood, abuse calcification of bones and muscle degeneration. The amount of lysine has increased in the cookies «Dachne» by 67 %, in the cookies «Yasne sonechko» by 68 % as compared with the check sample. The content of valine has increased in the cookies «Dachne» and «Yasne sonechko» by 2 %. The content of isoleucine has increased in the cookies «Dachne» by 21 % and «Yasne sonechko» by 15 %.

The biological value of food protein is characterized by the amino acid score; it is calculated as the ratio of the content of amino acids in the studied protein to their content in the conditionally ideal protein that meets the needs of the body. The amino acid score of new products is presented in Table 4. The increase of the lysine score in the new samples, especially in the «Yasne sonechko» is essential. The significant increase of the limited threonine score in the samples is a positive factor. The values of the score in both samples have exceeded 100 %.

Table 4
The amino acid score of the new sugar cookies, %

Amino acid	FAO/WHO scale, g/100 g of protein	Sample		
		Check sample	«Dachne»	«Yasne sonechko»
Isoleucine	4	65	66	67
Leucine	7	104	125	122
Lysine	5,5	47	66	68
Methionine + + cystine	3,5	71	77	78
Phenylalanine + + tyrosine	6	102	113	111
Threonine	4	85	100	102
Valine	5	67	57	59

The fraction of saturated fatty acids in all samples has been decreased as compared with the check sample and the fatty acid composition has been approached to "perfect lipid" owing to the usage of alternative oils in the production of new sugar cookies.

As seen from Fig. 2, the check sample is rich in saturated fatty acids (46, 95 %) and low in polyunsaturated fatty acids (13, 23 %). The fraction of fatty acids of cookies lipid base is lower (in the cookies «Dachne» by 14,65 %, in the cookies «Yasne sonechko» by 24,77 %) than it is in the check sample. Along with that, the fraction of polyunsaturated fatty acids is higher significantly.

For the normal functioning, the body needs biometals, currently divided into macro- and micronutrients. Due to the change of prescription, the mineral of sugar cookies has been improved (Table 5).




Fig. 2. Fatty acid composition of new samples

Table 6
Vitamin composition of the new cookies, mg %

Vitamin -	Check sample	«Dachne»	«Yasne sonechko»
Ascorbic acid	0,00	3,52±0,04	4,40±0,05
Thiamin	0,08±0,004	-	-
Tocopherol	0,26±0,002	-	9,73±0,60
Retinol	0,005±0,01	0,08±0,03	0,14±0,03

Note: «-» – not determined

The determination of the complex quality indicator has been carried out by organoleptic, physical and chemical, microbiological characteristics and the content of toxic elements, nutritional value, energetically value. The values of the complex quality indicator have been significantly increased in the new cookie samples (Fig. 3).

Table 5

The mineral composition of the new sugar cookies

№	Mineral elements	Check sample	«Dachne»	«Yasne sonechko»
Macroelements, mg				
1	Potassium (K)	163,02	247,03	308,97
2	Calcium (Ca)	90,00	278,00	206,00
3	Silicon (Si)	1,95	1,58	1,82
4	Magnesium (Mg)	14,75	22,56	26,80
5	Sodium (Na)	560,02	326,46	406,34
6	Sulphur (S)	45,42	38,86	42,61
7	Phosphorus (P)	260,00	406,00	380,00
Microelements, mkg				
8	Iron (Fe)	900,00	2500,00	1200,00
9	Iodine (I)	0,00	8,40	0,70
10	Manganese (Mn)	370,00	330,00	380,00
11	Selenium (Se)	5,92	3,69	4,71

The content of calcium has been increased in the new products by 2,28–3 times owing to fat-free whey powder and whey protein concentrate addition to the cookies recipes. The highest amount of it is achieved in the cookies "Dachne" due to the addition of the preparation of eggshell with lemon juice. The new products, especially the cookies "Dachne", have been enriched with iodine due to the changes in the recipe.

The study of the vitamin composition of sugar cookies (Table 6) shows that samples of cookies "Dachne" and "Yasne sonechko" are enriched with ascorbic acid, because it is absent in the check sample. The content of the tocopherol has been increased in the cookies "Yasne sonechko" by 37,4 times as compared with the check sample. Therefore, vitamin A content has been determined in the cookies "Yasne sonechko". Its amount has been increased by 28 times as compared with the check sample. It is also associated with the usage of processed products of sea buckthorn and calendula syrup and apricot powder in the recipe of cookies.




Fig. 3. The values of the complex quality indicators of the sugar cookies

Thus, the study of consumer properties of new types of cookies suggests that by organoleptic characteristics and nutritional value they exceed the check sample, it influenced the increase of the complex quality indicator. The physicochemical, toxicological and microbiological parameters meet the standard.

6. Discussing the research findings of consumer properties of the new sugar cookies

The results of the study, which are analyzed in the paper are characterized by combining the sea buckthorn jam with the calendula syrup in the fillings ("Yasne sonechko") and the usage of cherry plum and zucchini jam with the preparation of eggshell with lemon juice for the first time. The unconventional oils usage in the recipes has been also proposed.

Due to adjusting the recipes of sugar cookies, the consumer properties of the product have been improved. It is caused by the improvement of organoleptic parameters and increase of the beneficial nutritional compounds in the cookies composition. Satisfaction of the retinol daily needs by the intake of 100 grams of cookies "Dachne" has increased by 16 times compared to the check sample, through the usage of calendula powder and pumpkin oil. Satisfaction of the daily needs of calcium has increased (3 times higher than the check sample) and of the daily needs of iron has increased (2,7 times higher than check sample). The signifi-

cant increase of iodine in these cookies should also be noted. 100 g of the product by nearly 6 % satisfies the daily need of the substance. Satisfaction of the daily needs of fatty acids by consumption of 100 g of the product exceeds the similar parameter of the check sample by 2,4 times. The amount of amino acids in the cookies has increased by 1,8–3,5 times. In particular, it should be noted that the daily requirement for leucine and threonine by the intake of 100 grams of cookies is met by 79 and 80 % respectively. Cookies "Yasne sonechko" have been enriched with vitamins A, C and E. It can be explained by the usage of sea buckthorn oil and sea buckthorn jam in the cookies recipe. Due to the consumption of 100 grams of the new cookies, the daily needs of ascorbic acid are met by 6 %. The amount of vitamin A, as compared with the check sample, has increased by 28 times and the amount of vitamin E has increased by 37,4 times. The need for polyunsaturated fatty acids is satisfied almost by 91 %. It is 3,17 times higher than in the check sample. The amount of all essential acids has increased in the cookies and the lysine amount has increased most of all. Satisfaction of the daily needs of essential amino acids has increased by 1,67 times compared to the check sample.

However, along with the key benefits of the cookies, it should be noted that they are characterized by certain disadvantages. Thus, the usage of alternative raw materials, which has been proposed for the recipes of sugar cookies "Dachne" and "Yasne sonechko" is not usually available for the companies that produce large quantities of goods, because it requires additional resources. Also, non-traditional raw materials should be certified. It can also cause certain difficulties in the practical aspect. The usage of the additional raw materials along with traditional will influence the cost of products.

The results can be used for implementation in the confectionery companies of Ukraine and the European Union. Also, it can be used by other scientists because trends to reduce sugar and the total carbohydrate fraction are popular in the world. The developed fillings can be used for wafers, rolls, cakes and other confectionery products. The practical significance of the results is confirmed by patents of Ukraine for utility model "Composition of sugar cookies "Yasne sonechko" (№ 101331, 10.09.2015), "Composition of sugar cookies "Dachne" (№ 101329, 10.09.2015).

Further research is planned to devote to the study of changes in consumer properties during storage, the impact of packaging and different temperature regimes on the processes that occur during goods movement.

7. Conclusions

1. The recipes of new sugar cookies, which are characterized by the reduced content of wheat flour, sugar and fat have been developed. These components are replaced partly

by the unconventional raw materials. They contain a high amount of beneficial micronutrients. Thus, cookies "Dachne" also contain a vegetable and butter mixture, fat-free whey powder, whey protein concentrate, dried apple powder, dried raspberry leaf powder, calendula powder, pumpkin oil. Cherry plum and zucchini jam enriched with the preparation of eggshell with lemon juice has been used for the preparation of the filling. The cookies "Yasne sonechko" along with traditional raw materials contain whey protein concentrate, lungwort officinal powder, dried apricot powder, sea buckthorn oil. Sea buckthorn jam and calendula syrup have been used for the preparation of filling.

2. The new cookies samples have been evaluated by the parameters of taste and smell higher than the check sample especially according to the organoleptic evaluation. The cookies "Yasne sonechko" gained the most points. The developed samples meet the standard by the physical-chemical parameters and safety parameters. The energy value of the cookies "Dachne" and "Yasne sonechko" has reduced by 6 % and 3 % as compared with the check sample owing to partial replacement of high-calorie ingredients with unconventional raw materials. This factor has influenced the improvement of the nutritional value. Studies have shown that the reduction of fats and carbohydrates in all samples is achieved.

3. The determination of the fatty acid composition of sugar cookies allows to assert that the amount of saturated fatty acids decreased in all samples (in the cookies "Dachne" by 4,65 %, in the cookies "Yasne sonechko" by 24,77 %) compared to the check sample. Also, the amount of unsaturated fatty acids has increased in both samples by 20,03 and 29,67 times. These positive changes in the fatty acid composition of the products are due to the inclusion of vegetable pumpkin and sea buckthorn oils in the formulation. Adding of milk-containing products (whey protein concentrate, fat-free whey powder) and preparation of shell eggs has an impact on the increase in the amount of amino acids and improvement of the biological value of proteins. In particular, the amount of essential amino acids in the cookies "Dachne" has increased by 34,83 %. It also managed to improve the essential amino acid composition and to increase their content in the cookies "Yasne sonechko" by 32,54 % compared to the control sample. The content of calcium in new products increased by 2,28–3,0 times, which is also associated with the use of milk-containing products. The chemical composition of the filling has influenced the increase of iron, ascorbic acid, tocopherol and vitamin A in the samples.

4. The cookies "Dachne" and "Yasne sonechko" have higher complex quality indicators (0,83 and 0,84 units) due to significant improvement of the nutritional value and organoleptic characteristic of the developed products as compared with the check sample. The check sample according to the results of the qualimetric assessment has had 0,77 units only.

References

- Galushko, O. S. The trends of development market of confectionery and features of transformation in the system of values it's participants [Text] / O. S. Galushko // Actualni problemi ekonomici. – 2009. – Vol. 8. – P. 17–25.
- Tkachenko, A. S. The improvement of consumer properties of sugar cookies [Text] / A. S. Tkachenko, I. V. Syrokhman // Kharchova nauka i tekhnolohiya. – 2015. – Vol. 3. – P. 82–87.
- Bodak, M. P. The usage of non-traditional raw-materials for bakery [Text] / M. P. Bodak // Visnik Lvivskoi comerciynoi akademii. – 2014. – Vol. 14. – P. 113–116.
- Hadeeva, S. O. The determination potential risks of technology of baked biscuit semi with dietary supplements [Text] / S. O. Hadeeva, K. V. Svidlo // Zbirnik naukovih prats' NTU "KhPI". – 2010. – Vol. 46. – P. 276–283.

5. Tkachenko, A. S. The improvement of fat-acid contents of sugar cookies [Text] / A. S. Tkachenko // Visnik Lvivskoi comerciynoi akademii. – 2015. – Vol. 15. – P. 114–119.
6. Tkachenko, A. S. The sugar cookies with enriched protein contents [Text] / A. S. Tkachenko // Torgivlia, comercia, marketing. – 2015. – Vol. 18. – P. 118–122.
7. Shemanskaia, E. I. The phospholipid fat-containing products of functional appointment [Text] / E. I. Shemanskaia, N. I. Oseiko // The food science. – 2012. – Vol. 1. – P. 28–30.
8. Sincik, M. Investigation of Correlation between Traits and Path Analysis of Confectionary Sunflower Genotypes [Text] / M. Sincik, A. Goksoy // Notulae Botanicae Horti Agrobotanici Cluj-Napoca. – 2014. – Vol. 42, Issue 1. – P. 227–231. doi: 10.15835/nbha4219429
9. Ajai, B. C. Improving Genetic Attributes of Confectionary Traits in Peanut (*Arachis hypogaea L.*) Using Multivariate Analytical Tools [Text] / B. C. Ajay, M. V. C. Gowda, A. L. Rathnakumar, V. P. Kusuma, R. A. Fiyaz, P. Holajjer et. al. // Journal of Agricultural Science. – 2011. – Vol. 4, Issue 3. – P. 247–258. doi: 10.5539/jas.v4n3p247 .
10. Kulichenko, A. I. The usage of products with a milk raw material for confectionery products [Text] / A. I. Kulichenko // Molodoy uchenyy. – 2013. – Vol. 4. – P. 675–577.
11. Nikiforova, T. A. The prospects of usage secondary raw-materials of cereals productions [Text] / T. A. Nikiforova, S. M. Sevirenko, D. A. Kulikov et. al. // Khleboprodukti. – 2009. – Vol. 7. – P. 50–51.
12. Izembayeva Assel, K. Non-Traditional Raw Materials in Production of Sugar Cookies [Text] / Assel K. Izembayeva, Meruert P. Bayisbayeva, Bayan Z. H. Muldabekova et. al. // American–Eurasian J. Agric. & Environ. Sci. – 2014. – Vol. 3. – P. 358–362.
13. Maximova, A. A. Innovation technology of production of the oat cookies [Text] / A. A. Maximova // Khleboprodukti. – 2010. – Vol. 7. – P. 38–39.
14. Eckerson, J. M. The Effect of a Protein-Rich Egg White Breakfast and an Isocaloric Toaster Pastry Breakfast on Satiety and Dietary Intake in Normal-Weight College-Age Women [Text] / J. M. Eckerson, A. E. Lieb, K. A. Mullen, N. O. Allen // Journal of the Academy of Nutrition and Dietetics. – 2015. – Vol. 115, Issue 9 – P. A17. doi: 10.1016/j.jand.2015.06.048
15. Rosliakov, I. F. The usage of the Siberian pine nut kernel (*pinus sibirica*) in the production of pastry for functional appointment [Text] / I. F. Rosliakov, V. V. Gonchar, I. V. Shulvinskaia // Fundamental research. – 2007. – Vol. 7. – P. 89–90.
16. Rubilar, M. Flaxseed as a source of functional ingredients [Text] / M. Rubilar, C. Guti rrez, M. Verdugo, C. Shene, J. Sineiro // Journal of soil science and plant nutrition. – 2010. – Vol. 10, Issue 3. – P. 373–377. doi: 10.4067/s0718-95162010000100010
17. Li, S.-S. Fatty acid composition of developing tree peony (*Paeonia section Moutan DC.*) seeds and transcriptome analysis during seed development [Text] / S.-S. Li, L.-S. Wang, Q.-Y. Shu, J. Wu, L.-G. Chen, S. Shao, D.-D. Yin // BMC Genomics. – 2015. – Vol. 16, Issue 1. doi: 10.1186/s12864-015-1429-0
18. Fatima, T. Fatty Acid Composition of Developing Sea Buckthorn (*Hippophae rhamnoidesL.*) Berry and the Transcriptome of the Mature Seed [Text] / T. Fatima, C. L. Snyder, W. R. Schroeder, D. Cram, R. Datla, D. Wishart et. al. // PLoS One. – 2012. – Vol. 7, Issue 4. – P. e34099. doi: 10.1371/journal.pone.0034099
19. Dorokhovich, A. M. Fortification of the confectionery [Text] / A. M. Dorokhovich, O. L. Soloviova, V. V. Dorokhovich // Produkty & ingrediyenty. – 2011. – Vol. 3. – P. 26–28.
20. Bogatirov, A. Scientific principles of fortification nutrition products [Text] / A. Bogatirov // Khlibopekarska i kondyterska promyslovist Ukrayiny. – 2011. – Vol. 5. – P. 44–47.
21. Magomedov, G. O. Beet feelings for bakery [Text] / G. O. Magomedov, I. V. Plotnikova, N. G. Magomedov // Khleboprodukty. – 2014. – Vol. 8. – P. 44–47.
22. Alekseenko, E. V. The usage of see button concentrate in the cakes products [Text] / E. V. Alekseenko, I. M. Dikareva // Konditer-skoe i khlebopiekarskoe proizvodstvo. – 2014. – Vol. 5-6. – P. 6–8.
23. Ivanova, V. D. The researching of the cake functional properties [Text] / V. D. Ivanova, M. S. Khlebutina, N. P. Ivchuk // Naukovyi pratsi ONAKHT. – 2011. – Vol. 40, Issue 1. – P. 82–86.
24. Poliakova, A. V. Technology of the puff pastry with berries dry-powders supplements for puff pastry [Text] / A. V. Poliakova // Visnyk Donetskoho natsionalnoho universytetu ekonomiky i torhivli imeni Mykhayla Tuhan–Baranovskoho. Ser.: Tekhnichni nauky. – 2011. – Vol. 1. – P. 55–60.
25. Pop, A. Study of Valorification of *Lycium barbarum* (Goji) in Pastry Products [Text] / A. Pop, S. Muste, S. Man et. al. // Bulletin UASVM Food Science and Technology. – 2013. – Vol. 70, Issue 2. – P. 93–98.
26. Zucco, F. Physical and nutritional evaluation of wheat cookies supplemented with pulse flours of different particle sizes [Text] / F. Zucco, Y. Borsuk, S. Arnifield // LWT – Food Science and Technology. – 2012. – Vol. 44, Issue 10. – P. 2070–2076. doi: 10.1016/j.lwt.2011.06.007
27. Perez, S. Effect of soy flour and whey protein concentrate on cookie color [Text] / S. Perez, E. Matta, C. Osella, M. de la Torre, H. D. Sánchez // LWT – Food Sci. and Technol. – 2013. – Vol. 50, Issue 1. – P. 120–125. doi: 10.1016/j.lwt.2012.06.015
28. Gedrovica, P. Influence of Jerusalem Artichoke Powder on the Nutritional Value of Pastry Products [Text] / P. Gedrovica, D. Karklinia // International Journal of Biological, Biomolecular, Agricultural, Food and Biotechnological Engineering. – 2013. – Vol. 6. – P. 7.
29. Margie, M. C. Product Development of Malunggay (*Moringga Oleifera*) and Sweet Potato (*Ipomea Batatas*) for Pastry and Fillings [Text] / M. C. Margie, M. D. Manolito, P. R. L. Roey // Tropical Technology Journal. – 2012. – Vol. 19, Issue 1. – P. 5. doi: 10.7603/s40934-015-0005-1
30. Recipes of the cookie [Text]. – Moscow: Vsesoyuznyi nauchno-issledovatel'skiy institut konditerskoy promyshlenosti, 1987. – 248 p.

31. Krutoviy, Z. A. Matematichne modelyuvannya retsepturnoyi kompoztsiyi keksu pidvyshchenoyi kharchovoyi tsinnosti [Text] / Z. A. Krutoviy, G. V. Zakharenko, L. O. Kasilova et. al. // Nauka ta innovatsiyi. – 2013. – Vol. 5, Issue 9. – P. 5–9.
32. Regulation (EU) № 854 of the European Parliament and of the Council. Laying down specific rules for the organisation of official controls on products of animal origin intended for human consumption [Text]. – Official Journal of the European Union, 2004. – 45 p.
33. Ovchinnikova, I. A. New methods of analysis of amino acids, proteins and peptydov [Text] / I. A. Ovchinnikova. – Moscow: Myr, 1974. – 120 p.
34. Otsenka nekotorykh pyshchevykh dobavok [Text]. – 37-y doklad Obedynennoho komyteta ekspertov FAO/VOZ po pyshchevym dobavkam. Serrya tekhnicheskikh dokladov VOZ: Geneva, 1994. – P. 27–39.
35. AACC International. Methods 10-50D and 10-52. Approved Methods of the American Association of Cereal Chemists, 10th ed. [Text]. – AACC International, St. Paul, MN, U.S.A., 2000.
36. Tkachenko, A. S. The forecasting of demand on the new sugar cookie with non-traditional raw-materials [Text] / A. S. Tkachenko // Materialy IV regionalnoi konferencii: The goods of Lviv region on the today market: positioning, quality, safety within Eu-rointegration, 2015. – P. 220–224.

Досліджено можливість використання порошку з виноградних кісточок у технології здобного печива підвищеної біологічної цінності. Показано, що у порівнянні з пшеничним борошном він характеризується більшою водопоглинальною здатністю. Встановлено, що у присутності порошку з виноградних кісточок клейковина стає менш розтяжною та більш пружною. Доведено позитивний вплив порошку з виноградних кісточок на фізико-хімічні та органолептичні показники якості здобного печива

Ключові слова: виноградні вичавки, виноградні кісточки, порошок, клейковина, водопоглинальна здатність, здобне печиво

Исследована возможность использования порошка из виноградных косточек в технологии сдобного печенья повышенной биологической ценности. Показано, что по сравнению с пшеничной мукой он характеризуется большей водопоглотительной способностью. Установлено, что в присутствии порошка из виноградных косточек клейковина становится менее растяжимой и более упругой. Доказано положительное влияние порошка из виноградных косточек на физико-химические и органолептические показатели качества сдобного печенья

Ключевые слова: виноградные выжимки, виноградные косточки, порошок, клейковина, водопоглотительная способность, сдобное печенье

UDC 634.8: 664.682/.684

DOI: 10.15587/1729-4061.2016.69838

THE EFFECT OF GRAPE SEED POWDER ON THE QUALITY OF BUTTER BISCUITS

O. Samohvalova

PhD, Professor*

E-mail: sam-ov@mail.ru

N. Grevtseva

PhD, Associate Professor*

E-mail: nataver@yandex.ua

T. Brykova

Postgraduate Student*

E-mail: arizona19@mail.ru

A. Grigorenko

PhD, Chief technologist
Ltd "Charivna mozaika"

Avtostradna str., 21-A, Kharkiv, Ukraine, 61038

E-mail: anzhegrig@yandex.ru

*Department of Bakery, Confectionary,
Pasta and Food Concentrates Technology

Kharkiv State University of
Food Technology and Trade

Klochkivska str., 333, Kharkiv, Ukraine, 61051

1. Introduction

Governments of many countries are developing and implementing national programmes aimed at improvement of the nutritional status of population. These programmes include production of everyday food enriched with physiologically functional ingredients. In terms of environmental degradation and economic crisis, such products are essential for the health and working abilities of citizens and for providing a decent future via improved nutrition of children and youth.

All population groups are fond of pastries, in particular butter biscuits that are traditionally made of high-quality

wheat flour, margarine or other fats, sugar, and egg products. Although such biscuits contain much fat and a lot of carbohydrates and are devoid of biologically active substances. In this regard, introduction of useful ingredients – dietary fibre, polyphenolic compounds, minerals and vitamins – to the composition of biscuits has a beneficial effect on human health. Such components are abundant in raw plant materials, among which especially interesting is grape pomace (wine production waste). Ukraine is a wine-producing state due to favourable for growing grapes climatic conditions of southern regions and Transcarpathia. Annually, wineries of Ukraine accumulate large amounts of grape pomace after

ABSTRACT AND REFERENCES
TECHNOLOGY AND EQUIPMENT OF FOOD PRODUCTION

**RESEARCH OF SEDIMENTATION STABILITY
OF LIPID-MAGNETITE SUSPENSIONS BY THE
METHOD OF SPECTROPHOTOMETRY (p. 4-11)**

Alexandr Alexandrov, Iryna Tsykhanovska, Tatyana Gontar,
Nicholas Kokodiy

A spectrophotometric method of the assessment of stability and determining of the morphological characteristics of lipid-magnetite suspensions (LMS) was studied. The sizes of the particles of magnetite with a surface-active substance (SAS) were defined. The diameter of the particles is 78 nm. The concentration of the particles of magnetite stabilized by a surface-active substance was determined – the concentration (number in 1 cm³) equals N=1.33 · 10¹² cm⁻³ when obtaining a suspension. A slight decrease in the number of particles of magnetite with SAS in 1 cm³ of suspension was observed over time: during 48 hours, the concentration in 1 cm³ decreased from 1.33 · 10¹² down to 1.13 · 10¹² cm⁻³. The concentration decreases by approximately 2.25 % per 1 hour. Different LMS were obtained. The optimum ratios of the components were selected in the suspensions: magnetite, stabilizer and dispersion medium.

It was established that LMS can be used as biological-active additives, which possess comprehensive action: lipid-magnetite suspensions (LMS) on the basis of magnetite during oral introduction into human organism render beneficial biological effect with the period of action within the range of 3–4 hours: when entering LMS into human organism, the iron concentration in blood rises, which brings about:

- a short-term decrease in intracranial CSF pressure;
- activation of gastric and duodenum performance;
- increase in urination.

Due to bivalent iron and its ability to form transition complexes with oxygen and peroxide radicals (and hydro peroxides), magnetite also manifests antioxidant activity, which makes it possible to recommend it as an antioxidant, which facilitates improvement of the quality and prolongation of the period of storage of fat-containing products. Furthermore, LMS contain magnetite (which means digestible Fe²⁺); therefore they can be recommended as anti-anemic agent due to easily digestible bivalent iron. Thus, introduction of LMS into food products increases their quality, nutritional and biological value.

Therefore the studies of LMS are actual and they represent significant theoretical and practical interest.

Keywords: magnetite, suspension, method, SAS, sedimentation, stability, spectrophotometry, particle size, stabilization.

References

1. Chapa Gonzalez, C., Martínez Pérez, C. A., Martínez Martínez, A., Olivas Armendáriz, I., Zavala Tapia, O., Martel-Estrada, A., García-Casillas, P. E. (2014). Development of Antibody-Coated Magnetite Nanoparticles for Biomarker Immobilization. *Journal of Nanomaterials*, 2014, 1–7. doi: 10.1155/2014/978284
2. Unterweger, H., Tietze, R., Janko, C., Zaloga, J., Lyer, S., Taccardi, N. et. al. (2014). Development and characterization of magnetic iron oxide nanoparticles with a cisplatin-bearing polymer coating for targeted drug delivery. *International Journal of Nanomedicine*, 1, 3659–3676. doi: 10.2147/ijn.s63433
3. Cárdenas, W. H. Z., Mamani, J. B., Sibov, T., Caous, C. A., Amaro Jr., E., & Gamarrá, L. F. (2012). Particokinetics: computational analysis of the superparamagnetic iron oxide nanoparticles deposition process. *International Journal of Nanomedicine*, 2699–2712. doi: 10.2147/ijn.s30074
4. Cihanovskaja, I. V., Onoprienko, T. A., Kovalenko, V. A., Onoprienko, V. I. (2009). O perspektivah ispol'zovaniya magnetita v kachestve biologicheski aktivnykh dobavok. *Himija i tehnologija zhirov. Perspektivy razvitiya maslo-zhirovoj otrassli*. Kharkiv, NII masel i zhirov, 53–54.
5. Iljuha, N. G., Barsova, Z. V., Kovalenko, V. A., Cihanovskaja, I. V. (2010). Tehnologija proizvodstva i pokazateli kachestva pishhevoj dobavki na osnove magnetita. *Eastern-European Journal of Enterprise Technologies*, 6 (10 (48)), 32–35. Available at: <http://journals.uran.ua/eejet/article/view/5847/5271>
6. Denysova, A. Ju., Cyhanovskaja, Y. V., Skorodumova, O. B., Goncharenko, Ja. M., Pryjmak, G. O., Shevchenko, I. V. (2013). Doslidzhennja vplyvu zhyro-magnetytovoї suspensii' na termin zberigannja tvarynnyyh zhyriv. Part 1. Progresyvna tehnika ta tehnologii' harchovyh vyrobnyctv, restoranogo ta gotel'nogo gospodarstv i torgivli. Ekonomichna strategija i perspektyvy rozvytoku sfery torgivli ta poslug. Kharkiv, 71–72.
7. Cho, J., Koo, S. (2015). Characterization of particle aggregation in a colloidal suspension of magnetite particles. *Journal of Industrial and Engineering Chemistry*, 27, 218–222. doi: 10.1016/j.jiec.2014.12.038
8. Lou, W., Charalampopoulos, T. T. (1994). On the Electromagnetic Scattering and Ab-sorption of Agglomerated Small Spherical Particles. *Journal of Physics D: Applied Physics*, 27 (11), 2258–2270. doi: 10.1088/0022-3727/27/11/004
9. Xu, R. (2001). Particle Characterization: Light Scattering Methods. N.Y.: Kluwer Academic Publishers, 410.
10. Di Stasio, S. (2000). Feasibility of an optical experimental method for the sizing of primary spherules in sub-micron agglomerates by polarized light scattering. *Applied Physics B: Lasers and Optics*, 70 (4), 635–643. doi: 10.1007/s003400050872
11. Mulholland, G. W., Donnelly, M. K., Hagwood, C. R., Kukuck, S. R., Hackley, V. A., Pui, D. Y. H. (2006). Measurement of 100 nm and 60 nm Particle Standards by Differential Mobility Analysis. *Journal of Research of the National Institute of Standards and Technology*, 111 (4), 257–312. doi: 10.6028/jres.111.022
12. Ivanov, L. A., Kizevetter, D. V., Kiselev, N. N. et. al. (2006). Izmenenie svetovozvrashhenija ot stekljannyh mikrosharikov i prognoz kachestva sveto-vozvrashhajushhih pokrytij. *Opt. zhurn.*, 73 (1), 35–40.
13. Kizevetter, D. V., Maljugin, V. I. (2009). Odnovremennoe izmerenie razmerov i skorosti dvizhenija chastic. *Zhurn. tehn. fiziki*, 79 (2), 90–95.
14. Ershov, A. E., Isaev, I. L., Semina, P. N., Markel, V. A., Karpov, S. V. (2012). Effects of size polydispersity on the extinction spectra of colloidal nanoparticle aggregates. *Physical Review B*, 85 (4). doi: 10.1103/physrevb.85.045421
15. Karpov, S. V., Isaev, I. L., Gavriljuk, A. P. (2009). Opticheskie spektry kolloidov serebra s pozicijii fiziki fraktalov. *Kolloid. zhurn.*, 71 (3), 314.
16. Van de Hulst, G. (1961). *Rassejanie sveta malymi chasticami*. Moscow: IL, 536.
17. Kerker, M. (1969). The scattering of light and other electromagnetic radiation. N.Y., London, Academic Press, 666.
18. Xu, R. (2001). Particle Characterization: Light Scattering Methods. N.Y.: Kluwer Academic Publishers, 410.
19. Di Stasio, S. (2000). Feasibility of an optical experimental method for the sizing of primary spherules in sub-micron agglomerates by polarized light scattering. *Applied Physics B: Lasers and Optics*, 70 (4), 635–643. doi: 10.1007/s003400050872
20. Kizevetter, D. V., Maljugin, V. I. (2009). Odnovremennoe izmerenie razmerov i skorosti dvizhenija chastic. *Zhurn. tehn. fiziki*, 79 (2), 90–95.

21. Papok, I. M. (2012). Using the dynamic light-scattering method for the analysis of a blood-serum model solution. Moscow Univ. Phys. Bull., 67 (5), 452–456.
22. Iljuha, M. G., Cihanovs'ka, I. V., Barsova, Z. V., Timofeeva, V. P., Vedernikova, I. O. (2010). Patent. na korismu model' № 54284, MPK S 01 G 49/00. Sposob otrimannja magnetitu. Published 10.11.2010. Bull. № 21, 4.
23. Klenin, V. I., Shhegolev, S. Ju., Lavrushin, V. I. (1977). Haraktericheskie fu-nkciij svetorassejanija dispersnyh chastic. Saratov: Izd-vo SGU, 176.
24. Klenin, V. I. (1996). Praktikum po kolloidnoj himii. Moscow: Sol', 56.

DEVELOPING A MODEL OF THE FOAM EMULSION SYSTEM AND CONFIRMING THE ROLE OF THE YIELD STRESS SHEAR OF INTERFACIAL ADSORPTION LAYERS TO PROVIDE ITS FORMATION AND STABILITY (p. 11-19)

**Andrii Goralechuk, Svetlana Omel'chenko,
Oleg Kotlyar, Olga Grinchenko, Valery Mikhaylov**

The model of the formation of the foam emulsion by the emulsion whipping was developed. It was experimentally proved that the yield stress shear of interfacial adsorption layers can be used as a criterion for evaluating the stability of foams, emulsions and foam-emulsion systems. It was found that the introduction of DATEM to the reconstituted skimmed milk increases the yield stress shear of interfacial adsorption layers and stability of foams and emulsions. The introduction of lecithin's or DATEM reduces the yield stress shear of interfacial adsorption layers and stability of foams and emulsions accordingly. Simultaneous use of milk proteins, LACTEM, lecithin's and DATEM provides 1.3 times higher yield stress shear of interfacial adsorption layers at the water-air interface than at the water-oil interface, which is a thermodynamic condition for the formation of the foam emulsion by the emulsion whipping. It was proved that homogenization of the emulsion based on cocoa butter, milk proteins and surfactants provides destabilization of the emulsion and creates conditions for the flotation of destabilized fat particles.

The results allow justifying the parameters of the technology of the whipped semi-finished product based on cocoa butter, which is the emulsion whipping of which provides the foaming capacity of $450 \pm 22\%$, the mechanical strength of the foam emulsion of 3200 ± 160 Pa. It was confirmed that the whipping process can be divided into three stages: foaming, emulsion destabilization and adhesion of fat particles to air bubbles, providing high mechanical strength of the foam emulsion.

Keywords: interfacial adsorption layer, yield stress shear, whipped emulsion, foaming capacity.

References

1. Carr, N. O., Hogg, W. F. (2005). A manufacturer ' s perspective on selected palm-based products. Asia Pac J Clin Nutr, 14 (4), 381–386.
2. Allen, K. E., Murray, B. S., Dickinson, E. (2008). Development of a model whipped cream: Effects of emulsion droplet liquid/solid character and added hydrocolloid. Food Hydrocolloids, 22 (4), 690–699. doi: 10.1016/j.foodhyd.2007.01.017
3. Hotrum, N. E., Cohen Stuart, M. A., van Vliet, T., van Aken, G. A. (2004). Spreading of partially crystallized oil droplets on an air/water interface. Colloids and Surfaces A: Physicochemical and Engineering Aspects, 240 (1-3), 83–92. doi: 10.1016/s0927-7757(04)00172-4
4. Brun, M., Delample, M., Harte, E., Lecomte, S., Leal-Calderon, F. (2015). Stabilization of air bubbles in oil by surfactant crystals: A route to produce air-in-oil foams and air-in-oil-in-water emulsions. Food Research International, 67, 366–375. doi: 10.1016/j.foodres.2014.11.044
5. Fainerman, V. B., Aksenenko, E. V., Lylyk, S. V., Lotfi, M., Miller, R. (2015). Adsorption of Proteins at the Solution/Air Interface Influenced by Added Nonionic Surfactants at Very Low Concentrations for Both Components. 3. Dilational Surface Rheology. J. Phys. Chem. B, 119 (9), 3768–3775. doi: 10.1021/acs.jpcb.5b00136
6. Burke, J., Cox, A., Petkov, J., Murray, B. S. (2014). Interfacial rheology and stability of air bubbles stabilized by mixtures of hydrophobin and β -casein. Food Hydrocolloids, 34, 119–127. doi: 10.1016/j.foodhyd.2012.11.026
7. López-Castejón, M. L., de la Fuente, J., Ruiz, M., Guerrero, A. (2012). Influence of the presence of monoglyceride on the interfacial properties of soy protein isolate. Journal of the Science of Food and Agriculture, 92 (13), 2618–2623. doi: 10.1002/jsfa.5674
8. Fredrick, E., Heyman, B., Moens, K., Fischer, S., Verwijlen, T., Moldenaers, P. et al. (2013). Monoacylglycerols in dairy recombined cream: II. The effect on partial coalescence and whipping properties. Food Research International, 51 (2), 936–945. doi: 10.1016/j.foodres.2013.02.006
9. Phan, T. T. Q., Asaduzzaman, M., Le, T. T., Fredrick, E., Van der Meeren, P., Dewettinck, K. (2013). Composition and emulsifying properties of a milk fat globule membrane enriched material. International Dairy Journal, 29 (2), 99–106. doi: 10.1016/j.idairyj.2012.10.014
10. Kotlyar, O., Goralechuk, A., Grinchenko, O. (2014). The Study of Surface-Active Agents' Impact on the Strength of Interfacial Adsorption Layers. The Advanced Science Journal, 2014 (10), 37–42. doi: 10.15550/asj.2014.10.037
11. Eisner, M. D., Jeelani, S. A. K., Bernhard, L., Windhab, E. J. (2007). Stability of foams containing proteins, fat particles and nonionic surfactants. Chemical Engineering Science, 62 (7), 1974–1987. doi: 10.1016/j.ces.2006.12.056
12. Petkov, J. T., Gurkov, T. D., Campbell, B. E. (2001). Measurement of the Yield Stress of Gellike Protein Layers on Liquid Surfaces by Means of an Attached Particle. Langmuir, 17 (15), 4556–4563. doi: 10.1021/la001347i
13. Karbaschi, M., Lotfi, M., Kr gel, J., Javadi, A., Bastani, D., Miller, R. (2014). Rheology of interfacial layers. Current Opinion in Colloid & Interface Science, 19 (6), 514–519. doi: 10.1016/j.cocis.2014.08.003
14. Langevin, D. (2000). Influence of interfacial rheology on foam and emulsion properties. Advances in Colloid and Interface Science, 88 (1-2), 209–222. doi: 10.1016/s0001-8686(00)00045-2
15. Langevin, D., Marquez-Beltran, C., Delacotte, J. (2011). Surface force measurements on freely suspended liquid films. Advances in Colloid and Interface Science, 168 (1-2), 124–134. doi: 10.1016/j.cis.2011.03.007
16. Santini, E., Ravera, F., Ferrari, M., Stubenrauch, C., Makievski, A., Krägel, J. (2007). A surface rheological study of non-ionic surfactants at the water-air interface and the stability of the corresponding thin foam films. Colloids and Surfaces A: Physicochemical and Engineering Aspects, 298 (1-2), 12–21. doi: 10.1016/j.colsurfa.2006.12.004
17. Lexis, M., Willenbacher, N. (2014). Yield stress and elasticity of aqueous foams from protein and surfactant solutions – The role of continuous phase viscosity and interfacial properties. Colloids and Surfaces A: Physicochemical and Engineering Aspects, 459, 177–185. doi: 10.1016/j.colsurfa.2014.06.030
18. Maldonado-Valderrama, J., Martín-Rodriguez, A., Gálvez-Ruiz, M. J., Miller, R., Langevin, D., Cabrerizo-Vilchez, M. A. (2008). Foams and emulsions of β -casein examined by interfacial rheology. Colloids and Surfaces A: Physicochemical and Engineering Aspects, 323 (1-3), 116–122. doi: 10.1016/j.colsurfa.2007.11.003
19. Aksenenko, E. V., Kovalchuk, V. I., Fainerman, V. B., Miller, R. (2006). Surface dilational rheology of mixed adsorption layers at liq-

- uid interfaces. *Advances in Colloid and Interface Science*, 122 (1-3), 57–66. doi: 10.1016/j.cis.2006.06.012
20. Derkach, S. R., Krägel, J., Miller, R. (2009). Methods of measuring rheological properties of interfacial layers (Experimental methods of 2D rheology). *Colloid Journal*, 71 (1), 1–17. doi: 10.1134/s1061933x09010013
 21. Pelipenko, J., Kristl, J., Rošic, R., Baumgartner, S., Kocbek, P. (2012). Interfacial rheology: An overview of measuring techniques and its role in dispersions and electrospinning. *Acta Pharmaceutica*, 62 (2), 123–140. doi: 10.2478/v10007-012-0018-x
 22. Danov, K. D., Kralchevsky, P. A., Radulova, G. M., Basheva, E. S., Stoyanov, S. D., Pelan, E. G. (2015). Shear rheology of mixed protein adsorption layers vs their structure studied by surface force measurements. *Advances in Colloid and Interface Science*, 222, 148–161. doi: 10.1016/j.cis.2014.04.009
 23. Martin, A., Bos, M., Cohen Stuart, M., van Vliet, T. (2002). Stress-Strain Curves of Adsorbed Protein Layers at the Air/Water Interface Measured with Surface Shear Rheology. *Langmuir*, 18 (4), 1238–1243. doi: 10.1021/la011176x
 24. Izmajlova, V. N., Jampol'skaja, G. P., Summ, B. D. (1988). *Poverhnostnye javlenija v belkovyh sistemah*. Moscow: Himija, 240. Available at: <https://books.google.com.ua/books?id=rPOWtwAACAAJ>
 25. Denkov, N. D., Marinova, K. G., Tcholakova, S. S. (2014). Mechanistic understanding of the modes of action of foam control agents. *Advances in Colloid and Interface Science*, 206, 57–67. doi: 10.1016/j.cis.2013.08.004
 26. Omelchenko, S., Horalchuk, A., Hrynenko, O. (2014). Argumentation of Emulsifier Part in the Recipe of Foam and Emulsion Dairy Products Containing Vegetable Fats. *The Advanced Science Journal*, 2014 (7), 28–32. doi: 10.15550/asj.2014.07.028

OPTIMIZATION OF FORMULATION COMPOSITION OF THE LOW-CALORIE EMULSION FAT SYSTEMS (p. 20-27)

Nataliia Tkachenko, Pavlo Nekrasov,
Tetiana Makovska, Lubov Lanzhenko

Among the promising food products of oil-and-fat industry, special place is held by emulsion fat products, in which vegetable oil is in the dispersed state that increases its assimilation. High taste and nutritional properties, due to the specific character of their structure, are inherent in emulsion fat systems. Therefore, water-fat emulsions are the promising systems, on the basis of which it is possible to create mayonnaises, sauces, dressings, oil pastes, spreads and other food products, including low fat ones, with the balanced composition and health-improving properties.

The work substantiated the optimum content of the concentrate of Jerusalem artichoke "Noteo" and the stabilizing system "Hamulsion QNA" – 10,06 and 0,42 % respectively, as the components of low calorie emulsion fat basis for the production of low fat oil-and-fat products.

It is shown that a low calorie emulsion fat system, produced with the use of raw components in the optimal ratio, possesses standardized physical-chemical and microbiological indicators, high organoleptic characteristics and can be used as the raw material for the production of low calorie mayonnaises, sauces and dressings for healthy nutrition.

The recommendations are provided regarding the design of the technologies of the two groups of low calorie mayonnaises, sauces, dressings, enriched with food fibers and prebiotics (or the complexes of synbiotics), on the basis of the developed emulsion fat systems.

Keywords: low calorie emulsion system, Jerusalem artichoke, viscosity, organoleptic indicators, optimization, the response surface.

References

1. Smoljar, V. I. (2012). Stan faktychnogho kharchuvannja naselennja nezalezhnoji Ukrayiny. *Problemy kharchuvannja*, 1-2, 5–9.
2. Betoret, E., Betoret, N., Vidal, D., Fito, P. (2011). Functional foods development: Trends and technologies. *Trends in Food Science & Technology*, 22 (9), 498–508. doi: 10.1016/j.tifs.2011.05.004
3. Fahimdanesh, M., Mohammadi, N., Ahari, H., Khosravi, M. A., Zanjani, F. Zh. et. al. (2012). Effect of microencapsulation plus resistant starch on survival of *Lactobacillus casei* and *Bifidobacterium bifidum* in mayonnaise sauce. *African Journal of Microbiology Research*, 6 (40), 6853–6858. doi: 10.5897/ajmr12.1240
4. Mardar, M. R., Tkachenko, N. A., Lozovs'ka, Gh. M., Makovs'ka, T. V. (2016). Marketynghovi doslidzhennja pry pozycionuvanni ta vyvedenni na rynok nyzjokalorijnogho majonezu, zbaghachennogho kompleksom synbiotykiv. *Kharchova nauka i tekhnologija*, 1 (26), 3–10.
5. Ghazaei, S., Mizani, M., Piravi-Vanak, Z., Alimi, M. (2015). Particle size and cholesterol content of a mayonnaise formulated by OSA-modified potato starch. *Food Science and Technology (Campinas)*, 35 (1), 150–156. doi: 10.1590/1678-457x.6555
6. Kuo, S.-M., Merhige, P. M., Hagey, L. R. (2013). The Effect of Dietary Prebiotics and Probiotics on Body Weight, Large Intestine Indices, and Fecal Bile Acid Profile in Wild Type and IL10-/– Mice. *PLoS ONE*, 8 (3), e60270. doi: 10.1371/journal.pone.0060270
7. Harris, W. S., Miller, M., Tighe, A. P., Davidson, M. H., Schaefer, E. J. (2008). Omega-3 fatty acids and coronary heart disease risk: Clinical and mechanistic perspectives. *Atherosclerosis*, 197 (1), 12–24. doi: 10.1016/j.atherosclerosis.2007.11.008
8. Tkachenko, N. A., Sevast'janova, O. V., Makovs'ka, T. V. (2016) Zhyrozaminnyky vughlevodnoji ta bilkovoji pryrody v nyzjokalorijnykh majonezakh. *Prodovoljcha industrija APK*, 1-2, 18–22.
9. Mitchell, C. M., Davy, B. M., Halliday, T. M., Hulver, M. W., Neilson, A. P., Ponder, M. A., Davy, K. P. (2015). The effect of prebiotic supplementation with inulin on cardiometabolic health: Rationale, design, and methods of a controlled feeding efficacy trial in adults at risk of type 2 diabetes. *Contemporary Clinical Trials*, 45, 328–337. doi: 10.1016/j.cct.2015.10.012
10. Azizi, A., Homayouni, A., Payahoo, L. (2012). Effects of Probiotics on Lipid Profile: A Review. *American Journal of Food Technology*, 7 (5), 251–265. doi: 10.3923/ajft.2012.251.265
11. Mogilvii, M. P., Shaltumaev, T. Sh., Galukova, M. K. (2013). Sovremenniie napravleniya ispolzovaniia pishchevih volokon v kachestve funktsionalnih ingredientov. *Novie tekhnologii*, 1, 27–31.
12. Han, G. K. & Ko. (2006). Istorija uspeha. *Stabilizacionniie sistemi dlja idealnogo produkta. Pishchevaya promishlennost*, 8, 34.
13. Chatsisvili, N. T., Amvrosiadis, I., Kiosseoglou, V. (2012). Physico-chemical properties of a dressing-type o/w emulsion as influenced by orange pulp fiber incorporation. *LWT – Food Science and Technology*, 46 (1), 335–340. doi: 10.1016/j.lwt.2011.08.019
14. Myers, R., Montgomery, D., Anderson-Cook, C. (2016). Response surface methodology: process and product optimization using designed experiments. Hoboken, New Jersey: John Wiley & Sons, 825.
15. Tkachenko, N. A., Makovs'ka, T. V. (2015) Low-calorie mayonnaise production technology enriched with symbiotic complex by using batch method. *Journal of Food Science and Technology*, 4 (33), 74–81. doi: 10.15673/2073-8684.4/2015.55876
16. Rahbari, M., Aalam, M., Kashaninejad, M., Maghsoudlou, Y., Aghaei, S. S. A. (2014). A mixture design approach to optimizing low cholesterol mayonnaise formulation prepared with wheat germ protein isolate. *Journal of Food Science and Technology*, 52 (6), 3383–3393 doi: 10.1007/s13197-014-1389-4
17. Kishk, Y. F. M., Elsheshetawy, H. E. (2013). Effect of ginger powder on the mayonnaise oxidative stability, rheological measurements,

- and sensory characteristics. *Annals of Agricultural Sciences*, 58 (2), 213–220. doi: 10.1016/j.aoas.2013.07.016
18. Mardar, M. R., Valevsjka, L. O. (2010). Kompleksna tovaroznavchava ocinka jakosti novykh vydiv ekstrudovanykh zernovykh produktiv pidvyshhenoji kharchovoji cinnosti. *Zernovi produkty i kom-bikormy*, 1 (37), 19–22.
 19. Makovs'jka, T. V., Tkachenko, N. A. (2015). Bifidobacterium activation in technologies of health-improving mayonnaise. *Technology audit and production reserves*, 6 (4(26)), 40–44. doi: 10.15587/2312-8372.2015.56209

TECHNOLOGY OF SPECIALTY FATS BASED ON PALM STEARIN (p. 27-33)

Ekaterina Kunitsa, Aleksey Udovenko, Elena Litvinenko, Fedor Gladkiy, Irina Levchuk

New technology of modification of fats was developed, which allows, by fermentative ethanolysis, obtaining a new type of specialty fats for use in the food industry (culinary, bakery and dairy products). We proposed, for the modification of fatty raw materials, restructuring of the fats, namely, their active parts (acyl groups) with obtaining of derivatives of fatty acids that have functional properties. Obtained fats meet the requirements of normative documentation by the indicators of quality, and are additionally enriched with physiologically-active ingredients – ethyl esters of fatty acids, which are better digested and reduce the resynthesis of fat in a human body. The influence of conditions of fermentative alcoholysis of palm stearin by ethyl alcohol on the degree of its conversion to ethyl esters of fatty acids was defined.

It was established that when using ethyl alcohol as a reagent in the presence of lipolytic enzyme, ethyl esters and incomplete acylglycerols accumulate that causes the change of physical and chemical indicators (including the melting temperature decrease), the composition of the reaction mixture, and allows obtaining fats with given composition and properties. Thus, using this method, selecting necessary raw materials and varying the conditions of the reaction, one can obtain a whole range of specialty fats.

Keywords: alcoholysis, palm stearin, ethyl alcohol, enzyme, ethyl esters, specialty fat.

References

1. Sultanovich, Yu. A., Duxu, T. A. (2015). Problemy pry prymenenyye bestransovyyh zhryrov v kondyterskom proyzvodstve. *Masla y zhryry*, 3-4, 23–25.
2. Zajceva, L. V. (2015). Trans-zyomery zhrynyh kyslot – vred y opasnost. *Masla y zhryry*, 3-4, 25–27.
3. Timms, R. E. (2005). Fractional crystallisation - the fat modification process for the 21st century. *European Journal of Lipid Science and Technology*, 107 (1), 48–57. doi: 10.1002/ejlt.200401075
4. Noipin, K., Kumar, S. (2014). Optimization of ethyl ester production from palm oil. *Petroleum & Coal*, 56 (3), 249–258.
5. Korus, R. A., Hoffman, D. S., Bam, N., Peterson, C. L., Drown, D. C. (1995). Transesterification process to manufacture ethyl ester of rape oil. *First Biomass Conference of the Americas: Energy, Environment, Agriculture, and Industry*, 2, 815–826.
6. Latip, R. A., Lee, Y.-Y., Tang, T.-K., Phuah, E., Tan, C., Lai, O.-M. (2013). Physicochemical properties and crystallisation behaviour of bakery shortening produced from stearin fraction of palm-based diacylglycerol blended with various vegetable oils. *Food Chemistry*, 141 (4), 3938–3946.
7. Abd. Rashid, N., Chiew Let, C., Chong Seng, C., Omar, Z. (2012). Crystallisation kinetics of palm stearin, palm kernel olein and their blends. *LWT - Food Science and Technology*, 46 (2), 571–573. doi: 10.1016/j.lwt.2011.11.001
8. Barison, Dr. Y., Sundram, Dr. K. (Eds.) (2014). *Palmovoe maslo Malajzii. Obogashchaya zhyn. Dnepropetrovsk, Ekspert Agro*, 44.
9. Nekrasov, P. A., Podlynsa, O. V., Gopkalov, V. G. (2014). Yssledovaniye pyshhevoj cennosty dyacylglycerynovogo masla. *Masla y zhryry*, 3-4, 7–9.
10. Xarkevych, D. A. (2006). *Farmokologyya*. Moscow: GEOTAR-Meda, 560.
11. Korenskaya, Y. M., Yvanovskaya, N. P., Kolosova, O. A. (2008). Lekarstvennye rastenyya y lekarstvennoe rastytelnoe syre, soderzhashhye vytamyny, polysaxarydy, zhyrnye masla. Voronezh, Yzatelsko-polygraficheskij centr Voronezhskogo gosudarstvennogo unversyteta, 320.
12. Kaprelyancz, L. V., Iorgachova, K. G. (2003). *Funkcionalni produkty*. Odesa: Druk, 312.
13. Laposata, M. (1999). Fatty acid ethyl esters: Nonoxidative ethanol metabolites with emerging biological and clinical significance. *Lipids*, 34 (S1), 281–285. doi: 10.1007/bf02562318
14. Tereshhuk, L. V., Mamontov, A. S., Starovojtova, K. V. (2014). Produkty frakcyonirovannyya palmovogo masla v proyzvodstve spredov. *Texnika y texnologyya pyshhevyx proyzvodstv*, 3, 79–83.
15. Tyutyunnikov, B. N., Buhshtab, Z. I., Gladkij F. F. et al. (1992). *Himija zhirov*. Moscow: Kolos, 448.
16. Garabadzhyu, A. V., Galynkyn, V. A., Karasev, M. M., Kozlov, G. V., Lysyczkaya, T. V. (2010). Osnovnye aspektы yspolzovaniyya lypaz dlya poluchenyya byodyzelya. *Yzvestyya Sankt-Peterburgskogo gosudarstvennogo texnologicheskogo yn-ta (texnicheskogo un-ta)*, 7, 63–67.

THE INFLUENCE OF MECHANOLYSIS ON THE ACTIVATION OF NANOCOMPLEXES OF HETEROPOLYSACCHARIDES AND PROTEINS OF PLANT BIOSYSTEMS IN DEVELOPING OF NANOTECHNOLOGIES (p. 33-40)

Raisa Pavlyuk, Viktoriya Pogarska, Tatyana Kotuyk, Aleksey Pogarskiy, Svitlana Loseva

A nanotechnology of protein plant supplements in the form of puree of peas was developed that is based on the processes of deep processing of raw materials. Finely dispersed grinding and steam and thermal processing were used in this work as the innovation. When using traditional methods of raw materials processing, biological potential is not used in full.

It was found that during deep processing of plant raw materials (dried peas), which is based on comprehensive effect of steam and thermal processing and finely dispersed grinding on the raw material in obtaining nanostructured puree, the processes of mechanical destruction and mechanical chemistry occur. These processes are accompanied by non-enzymatic biocatalysis – mechanolysis (destruction) of hard soluble biopolymers and nanocomplexes of biopolymers (proteins, heteropolysaccharides, namely, pectins, cellulose, starch) with their transformation to monomers (35...55 %) into soluble easily absorbed form (almost 2 times higher compared to the original raw material in a hidden form). The mechanism of protein mechanical destruction and its nanocomplexes, which is associated with the mechanical cracking, was discovered. It was found that the steam and thermal processing and finely dispersed grinding of peas, while obtaining finely dispersed puree, leads to the destruction of polysaccharides by the non-enzymatic catalysis, namely cellulose and starch (30...35 %), protopectin (50 %), to separate monomers. It is shown that in parallel there is an increase of glucose in nanopuree of peas (1.0 g ...10.0 g/100 g, i. e. by 10 times).

Integrated application of these processes is accompanied by mechanical destruction, mechanical activation and mechanolysis of

biopolymers of nanocomplexes (protein, heteropolysaccharides, etc.) to α -amino acids, glucose, etc. (48...55 %).

Keywords: nanotechnologies, finely dispersed grinding, mechanoysis, nanocomplexes, biopolymers, heteropolysaccharides.

References

- FAO/WHO/UNU. Dietary protein quality evalution in human nutrition (2013). Report of an FAO Expert Consultation. Food and agriculture organization of the united nations Rome, 92, 57.
- Kaprel'janc, L. V. (2015). Prebiotiki: himija, tehnologija, primenenie. Kyiv: EnterPrint, 252.
- Gibson, G., Roberfroid, M. (2008). Handbook of Prebiotics. Vol. 4. CRS Press, London, 22–42.
- Sousa, V. M. C. de, Santos, E. F. dos, Sgarbieri, V. C. (2011). The Importance of Prebiotics in Functional Foods and Clinical Practice. *Food and Nutrition Sciences*, 02 (02), 133–144. doi: 10.4236/fns.2011.22019
- Roberfroid, M. B. (2000). Fructo-oligosaccharide malabsorption: benefit for gastrointestinal functions. *Current Opinion in Gastroenterology*, 16 (2), 173–177. doi: 10.1097/00001574-200003000-00013
- Pavljuk, R. Ju., Pogarskaja, V. V., Pavljuk, V. A., Radchenko, L. A., Jur'eva, O. A., Maksimova, N. F. (2015). Krio- i mehanohimija v pishhevyyh tehnologijah. Har'kovskij gosudarstvennyj universitet pitanija i torgovli; Har'kovskij torgovo-ekonomicheskij institut; Kievskij nacional'nyj torgovo-ekonomicheskij universitet, 255.
- Galland, L. (2014). Functional Foods: Health Effects and Clinical Applications. 3rd edition. Reference Module in Biomedical Sciences, from Encyclopedia of Human Nutrition, 366–371.
- Tur, J. A., Bibiloni, M. M. (2015). Functional Foods. Reference Module in Food Science. Encyclopedia of Food and Health, 157–161.
- Tu, J., Zhang, M., Xu, B., Liu, H. (2015). Effects of different freezing methods on the quality and microstructure of lotus (*Nelumbo nucifera*) root. *International Journal of Refrigeration*, 52, 59–65. doi: 10.1016/j.ijrefrig.2014.12.015
- Goñi, I., Serrano, J., Saura-Calixto, F. (2006). Bioaccessibility of β -Carotene, Lutein, and Lycopene from Fruits and Vegetables. *Journal of Agricultural and Food Chemistry*, 54 (15), 5382–5387. doi: 10.1021/jf0609835
- Bernstein, P. S., Khachik, F., Carvalho, L. S., Muir, G. J., Zhao, D.-Y., Katz, N. B. (2001). Identification and Quantitation of Carotenoids and their Metabolites in the Tissues of the Human Eye. *Experimental Eye Research*, 72 (3), 215–223. doi: 10.1006/exer.2000.0954
- Dherani, M., Murthy, G. V. S., Gupta, S. K., Young, I. S., Maraini, G., Camparini, M. et al. (2008). Blood Levels of Vitamin C, Carotenoids and Retinol Are Inversely Associated with Cataract in a North Indian Population. *Investigative Ophthalmology & Visual Science*, 49 (8), 3328–3335. doi: 10.1167/iovs.07-1202
- Goñi, I., Serrano, J., Saura-Calixto, F. (2006). Bioaccessibility of β -Carotene, Lutein, and Lycopene from Fruits and Vegetables. *Journal of Agricultural and Food Chemistry*, 54 (15), 5382–5387. doi: 10.1021/jf0609835
- Bernstein, P. S., Khachik, F., Carvalho, L. S., Muir, G. J., Zhao, D.-Y., Katz, N. B. (2001). Identification and Quantitation of Carotenoids and their Metabolites in the Tissues of the Human Eye. *Experimental Eye Research*, 72 (3), 215–223. doi: 10.1006/exer.2000.0954
- Gerasymenko, S. S., Gerasymenko, V. S. (2013). Statystychna harkterystyka spozhyvannja produktiv harchuvannja naselennjam Ukrai'ny. *Statystyka Ukrai'ny*, 2, 28–33.
- Dejneko, L. V., Shelud'ko, E. I. (2013). Harchova promyslovist' Ukrai'ny: efektyvnist' vykorystannja vyrobnychih resursiv ta kadrovo-goto potencialu. NAN Ukrai'ny, DU "In-t ekon. ta prognoz. NAN Ukrai'ny", 120.
- James, S. J., James, C. (2014). Chilling and Freezing. *Food Safety Management*, 481–510. doi: 10.1016/b978-0-12-381504-0.00020-2
- Shi, L., Li, W., Sun, J., Qiu, Y., Wei, X., Luan, G. et al. (2016). Grinding of maize: The effects of fine grinding on compositional, functional and physicochemical properties of maize flour. *Journal of Cereal Science*, 68, 25–30. doi: 10.1016/j.jcs.2015.11.004
- Balaz, P. (2010). Mechanochemistry in Nanoscience and Minerals Engineering. Woodhead Publishing Limited, 400.
- Baláž, P., Baláž, M., Bujňáková, Z. (2014). Mechanochemistry in Technology: From Minerals to Nanomaterials and Drugs. *Chemical Engineering & Technology*, 37 (5), 747–756. doi: 10.1002/ceat.201300669
- Boldyrev, V. V. (2004). Mechanochemical modification and synthesis of drugs. *Journal of Materials Science*, 39 (16/17), 5117–5120. doi: 10.1023/b:jmsc.0000039193.69784.1d

STUDYING THE ACCUMULATION OF NITROGENOUS SUBSTANCES IN BIOFORTIFIED PUMPKIN VEGETABLES (p. 40-46)

Gregoriy Deinychenko, Olha Yudicheva

The main purpose of biofortification is obtaining plant products with improved nutritional properties. Plant products are biofortified by means of the classic selection, genetic modification, or with the use of special fertilizers. Food plants have traditionally been enriched with vital minerals and vitamins; lately, they have also been bioenriched with amino acids and proteins. Vegetable protein consumed with the animal one enhances the value of protein nutrition due to the formed biologically active amino acid complexes. The value of vegetable protein increases in vegetarian nutrition, especially hard food, and nutrition of people suffering from celiac disease. We have studied the peculiarities of nitrogenous substances' accumulation in biofortified pumpkin vegetables grown with the use of the liquid, organic, environment-friendly Riverm fertilizer. The objects of study are biofortified pumpkin vegetables: pumpkins of Oleshkivskyi and Sviten varieties, melons of Olvia and Fortuna varieties, and watermelons of Orphei and Atlant varieties. The reference samples are vegetables grown by the standard technology, without the above mentioned fertilizer. The research findings show that biofortified pumpkin vegetables are characterized by higher contents of total nitrogen and protein nitrogen, as well as contain more protein in comparison with the reference samples: pumpkins – by 15.0–17.6 %, melons – by 6.5–16.4 %, and watermelons – by 8.9–10.1 %. The highest amount of essential amino acids is contained in the protein of biofortified pumpkins, a bit lower – in biofortified melons and watermelons. The protein of biofortified pumpkins is characterized by the content of leucine, valine, and lysine. Biofortified melons and watermelons are dominated by lysine and phenylalanine. The largest shares of replaceable amino acids in all the samples are those of aspartic acid and glutamic acid. Bioenriched with nitrogenous substances (in particular, protein and amino acids) pumpkin vegetables cannot fully satisfy human needs of proteins and essential amino acids, although they can perfectly supplement nutrition with the latter. Such vegetables can be recommended to be used in balanced diets of animal and vegetable proteins, glutenless diets, and vegetarian diets.

Keywords: biofortification, fertilizers, Riverm, protein, nitrogenous substances, amino acids, pumpkin vegetables, micronutrients.

References

- Dunaevskiy, G. A., Popik, S. Y. (1990). Ovoshchi i frukty v pitaniu zdorovogo i bolnogo cheloveka. Kyiv: Zdorove, 160.
- Hirsch, K. D. (2009). Nutrient Biofortification of Food Crops. *Annual Review of Nutrition*, 29 (1), 401–421. doi: 10.1146/annurev-nutr-080508-141143
- Burlaka, O. M., Sorochynskyy B. V. (2010) Roslynni biotekhnolohii: biofortifikatsiia kharchovykh roslyn. Kyiv: DIA, 88.

4. Welch, R. M. (2005). Biotechnology, Biofortification, and Global Health. *Food and Nutrition Bulletin*, 26 (4), 304–306. doi: 10.1177/15648265050264s309
5. Fageria, N. K., Moraes, M. E., Ferreira, E. P. B., Knupp, A. M. (2012). Biofortification of Trace Elements in Food Crops for Human Health. *Communications in Soil Science and Plant Analysis*, 43 (3), 556–570. doi: 10.1080/00103624.2012.639431
6. Murgia, I., De Gara, L., Grusak, M. A. (2013). Biofortification: how can we exploit plant science and biotechnology to reduce micronutrient deficiencies? *Frontiers in Plant Science*, 4, 429. doi: 10.3389/fpls.2013.00429
7. Chojnacka, K., Mikulewicz, M., Cieplik, J. (2011). Biofortification of Food with Microelements. *American Journal of Agricultural and Biological Sciences*, 6 (4), 544–548. doi: 10.3844/ajabssp.2011.544.548
8. Mayer, J. E., Pfeiffer, W. H., Beyer, P. (2008). Biofortified crops to alleviate micronutrient malnutrition. *Current Opinion in Plant Biology*, 11 (2), 166–170. doi: 10.1016/j.pbi.2008.01.007
9. Gilligan, D. O. (2012). Biofortification, Agricultural Technology Adoption, and Nutrition Policy: Some Lessons and Emerging Challenges*. *CESifo Economic Studies*, 58 (2), 405–421. doi: 10.1093/cses/ifs020
10. Leyva-Guerrero, E., Narayanan, N. N., Ihemere, U., Sayre, R. T. (2012). Iron and protein biofortification of cassava: lessons learned. *Current Opinion in Biotechnology*, 23 (2), 257–264. doi: 10.1016/j.copbio.2011.12.009
11. DellaValle, D. M., Thavarajah, D., Thavarajah, P., Vandenberg, A., Glahn, R. P. (2013). Lentil (*Lens culinaris* L.) as a candidate crop for iron biofortification: Is there genetic potential for iron bioavailability? *Field Crops Research*, 144, 119–125. doi: 10.1016/j.fcr.2013.01.002
12. Nair, R. M., Yang, R.-Y., Easdown, W. J., Thavarajah, D., Thavarajah, P., Hughes, J. d'A, Keatinge, J. D. (2013). Biofortification of mungbean (*Vigna radiata*) as a whole food to enhance human health. *Journal of the Science of Food and Agriculture*, 93 (8), 1805–1813. doi: 10.1002/jsfa.6110
13. McGrath, S. P., Chambers, B. J., Taylor, M. J., Carlton-Smith, C. H. (2012). Biofortification of zinc in wheat grain by the application of sewage sludge. *Plant and Soil*, 361 (1-2), 97–108. doi: 10.1007/s11104-012-1381-6
14. Aciksoz, S. B., Yazici, A., Ozturk, L., Cakmak, I. (2011). Biofortification of wheat with iron through soil and foliar application of nitrogen and iron fertilizers. *Plant and Soil*, 349 (1-2), 215–225. doi: 10.1007/s11104-011-0863-2
15. Hussain, S., Maqsood, M. A., Rengel, Z., Aziz, T. (2012). Biofortification and estimated human bioavailability of zinc in wheat grains as influenced by methods of zinc application. *Plant and Soil*, 361 (1-2), 279–290. doi: 10.1007/s11104-012-1217-4
16. Zhang, Y.-Q., Deng, Y., Chen, R.-Y., Cui, Z.-L., Chen, X.-P., Yost, R. et. al. (2012). The reduction in zinc concentration of wheat grain upon increased phosphorus-fertilization and its mitigation by foliar zinc application. *Plant and Soil*, 361 (1-2), 143–152. doi: 10.1007/s11104-012-1238-z
17. Ajiboye, B., Cakmak, I., Paterson, D., de Jonge, M. D., Howard, D. L., Stacey, S. P. et. al. (2015). X-ray fluorescence microscopy of zinc localization in wheat grains biofortified through foliar zinc applications at different growth stages under field conditions. *Plant and Soil*, 392 (1-2), 357–370. doi: 10.1007/s11104-015-2467-8
18. Zou, C. Q., Zhang, Y. Q., Rashid, A., Ram, H., Savasli, E., Ariyossy, R. Z. et. al. (2012). Biofortification of wheat with zinc through zinc fertilization in seven countries. *Plant and Soil*, 361 (1-2), 119–130. doi: 10.1007/s11104-012-1369-2
19. Poblaciones, M. J., Rodrigo, S. M., Santamaría, O. (2012). Evaluation of the Potential of Peas (*Pisum sativum* L.) to Be Used in Selenium Biofortification Programs Under Mediterranean Conditions. *Biological Trace Element Research*, 151 (1), 132–137. doi: 10.1007/s12011-012-9539-x
20. Rahmana, M. M., Erskine, W., Zaman, M. S., Thavarajah, P., Thavarajah, D., Siddique, K. H. M. (2013). Selenium biofortification in lentil (*Lens culinaris* Medikus subsp. *culinaris*): Farmers' field survey and genotype × environment effect. *Food Research International*, 54 (2), 1596–1604. doi: 10.1016/j.foodres.2013.09.008
21. Seppänen, M. M., Kontturi, J., Heras, I. L., Madrid, Y., Cámaras, C., Hartikainen, H. (2010). Agronomic biofortification of Brassica with selenium – enrichment of SeMet and its identification in Brassica seeds and meal. *Plant and Soil*, 337 (1-2), 273–283. doi: 10.1007/s11104-010-0523-y
22. Landini, M., Gonzali, S., Perata, P. (2011). Iodine biofortification in tomato. *Journal of Plant Nutrition and Soil Science*, 174 (3), 480–486. doi: 10.1002/jpln.201000395
23. Blasco, B., Ríos, J. J., Leyva, R., Cervilla, L. M., Sánchez-Rodríguez, E., Rubio-Wilhelmi, M. M. et. al. (2010). Does Iodine Biofortification Affect Oxidative Metabolism in Lettuce Plants? *Biological Trace Element Research*, 142 (3), 831–842. doi: 10.1007/s12011-010-8816-9
24. Voogt, W., Holwerda, H. T., Khodabaks, R. (2010). Biofortification of lettuce (*Lactuca sativa* L.) with iodine: the effect of iodine form and concentration in the nutrient solution on growth, development and iodine uptake of lettuce grown in water culture. *Journal of the Science of Food and Agriculture*, 90 (5), 906–913. doi: 10.1002/jsfa.3902
25. Jin, Z., Minyan, W., Lianghuan, W., Jiangguo, W., Chunhai, S. (2008). Impacts of Combination of Foliar Iron and Boron Application on Iron Biofortification and Nutritional Quality of Rice Grain. *Journal of Plant Nutrition*, 31 (9), 1599–1611. doi: 10.1080/01904160802244803
26. Ježek, P. et. al. (2011). Effect of foliar application of selenium on the content of selected amino acids in potato tubers (*Solanum tuberosum* L.). *Plant soil environ.*, 57 (7), 315–320.
27. Kozak, V. V. (2009). Printsipy ekologicheskogo bezopasnogo zemledeliia. Kyiv: MEF «AQUA-VITAE», 38.
28. Yudicheva, O. (2014). Study of Zinc Content in Biofortified Tomato. *The Advanced Science Journal*, 2014 (7), 15–18. doi: 10.15550/asj.2014.07.015
29. Dejnychenko, G. V., Judicewa, O. P. (2015). Study of possibilities to grow biofortified vegetables as a source of carotenoids. *Eastern-European Journal of Enterprise Technologies*, 2/10 (74), 36–40. doi: 10.15587/1729-4061.2015.39763
30. Yudicheva, O. P. (2015). Zavisimost khimicheskogo sostava ot sorta biofortifitsirovannykh tykvennykh ovoshchey. *Vestnik Sibirskogo universiteta potrebitelskoy kooperatsii*, 4 (11), 68–72.

RESEARCH INTO THE IMPACT OF ENZYME PREPARATIONS ON THE PROCESSES OF GRAIN DOUGH FERMENTATION AND BREAD QUALITY (p. 46-53)

Svitlana Oliinyk, Olga Samokhvalova, Anna Zaparenko, Elena Shidakova-Kamenyuka, Nicola Chekanov

The important issue of improvement in the technology of grain bread is the development of measures to improve the rheological properties of dough and bread. With this aim we proposed the use of cellulases, hemicellulases and oxidoreductases at the stage of dough mixing. It is shown that the application of the studied enzymes for grain emmer and wheat dough contributes to the intensification of non-starch polysaccharides hydrolysis, namely reduction of the content of cellulose by 11 %, hemicelluloses – by 14.3 and 13.0 %, and increase in the content of water-soluble fraction of hemicelluloses.

Additionally, the presence of enzyme preparations in the grain dough promotes slowing down of the processes of gluten proteolysis and improvement of its rheological properties that predetermines increase of gas-retaining capacity of the studied system. It was discovered that adding the studied enzyme preparations to grain dough contributes to the intensification of acid and gas generation in it. The resulting effect of biochemical and microbiological changes in grain dough under the influence of enzyme preparations of cellulase, xylanase and glucose oxidase is the improvement in the quality of grain bread compared to the samples without their addition.

Keywords: enzymes, grain bread, dough preparation, cellulase, xylanase, glucose oxidase.

References

1. Mofidi, A., Ferraro, Z. M., Stewart, K. A., Tulk, H. M. F., Robinson, L. E., Duncan, A. M., Graham, T. E. (2012). The Acute Impact of Ingestion of Sourdough and Whole-Grain Breads on Blood Glucose, Insulin, and Incretins in Overweight and Obese Men. *Journal of Nutrition and Metabolism*, 2012, 1–9. doi: 10.1155/2012/184710
2. Shkapov, E. I. (2002). Sovershenstvovaniye technologii dispergirovaniya zerna dlia proizvodstva khlebobulochnykh izdeliy. Moskovskij gosudarstvennyj universitet pishhevyh proizvodstv, 18.
3. Koriachkina, S. Ya., Kuznetsova, E. A., Cherepnina, L. V. (2012). Technologiya khleba iz tselogo zerna triticale. Orel: FGBOU VPO «Gosuniversitet – UNPK», 177.
4. Balanov, P. E., Smotrayeva, I. V. (2014). Technologiya soloda. SPb.: NIU ITMO; IHIBT, 82.
5. Pshenyshniuk, G. F., Makarova, O. V., Ivanova, G. S. (2010). Innovativnye zahody pidvyshchennia yakosti zernovogo khliba. Kharchova nauka i technologiya, 1, 73–77.
6. Koriachkina, S. Ya., Kuznetsova, E. A., Prigarina, O. M. (2006). Povysheniye mikrobiologicheskoy chistoty rzhyy i pshenitsy na stadii zamachivaniya v rastvorakh chimicheskikh sredstv pri proizvodstve zernovogo khleba. Prioritetnye i nauchnoye obespecheniye realizatsyi gosudarstvennoy politiki zdorovogo pataniya, 25–28.
7. Ivanova, G. S., Leschuk, O. V. (2012) Vplyv sposobiv tistoprygotuvannia na yakist zernovogo khliba na osnovi sumishi. Naukovy zdobutky molodi – vyrihenniu problem kharchuvannia liudstva u XXI st., 1, 88.
8. Novikova, A. N. (2004). Sovremennaya technologiya khleba iz tselogo zerna pshenitsy. Moskovskij gosudarstvennyj universitet pishhevyh proizvodstv, 25.
9. Kalina, M. A. (2012). Razrabotka zernovogo khleba iz triticale i otsenka yego potrebitelskikh svoystv. FGBOU VPO «Rossijskij ekonomicheskij universitet im. G. V. Plehanova», 22.
10. Makarova, O. V., Pshenyshniuk, G. F., Ivanova, G. S. (2011). Vliyanie vlagoteplovoy obrabotki pshenitsy na pokazateli kachestva zernovogo khleba. Kharchova nauka i technologiya, 1, 69–72.
11. Kaprelyants, L. V. (2009). Phermenty v piashhevyykh technologiyakh. Odessa, 468.
12. Melim Miguel, A. S., Souza, T., Costa Figueiredo, E. V. da, Paulo Lobo, B. W., Maria, G. (2013). Enzymes in Bakery: Current and Future Trends. *Food Industry*. doi: 10.5772/53168
13. Ognean, M., Ognean, C. F., Bucur, A. (2011). Rheological Effects of Some Xylanase on Doughs from High and Low Extraction Flours. *Procedia Food Science*, 1, 308–314. doi: 10.1016/j.profoo.2011.09.048
14. Shah, A. R., Shah, R. K., Madamwar, D. (2006). Improvement of the quality of whole wheat bread by supplementation of xylanase from *Aspergillus foetidus*. *Bioresource Technology*, 97 (16), 2047–2053. doi: 10.1016/j.biortech.2005.10.006
15. Sanina, T. V., Shuvayeva, G. P., Aliokhina, N. N. (2003). Intensifikatsiya protsessa bioaktivatsyi zerna i snizheniye yego mikrobiologicheskoy obsemennostti v technologii zernovogo khleba. *Hranenye i pererabotka sel'hozsyryja*, 1, 15–17.
16. Renzetti, S., Rosell, C. M. (2016). Role of enzymes in improving the functionality of proteins in non-wheat dough systems. *Journal of Cereal Science*, 67, 35–45. doi: 10.1016/j.jcs.2015.09.008
17. Bonet, A., Rosell, C. M., P rez-Munuera, I., Hernando, I. (2007). Rebuilding gluten network of damaged wheat by means of glucose oxidase treatment. *Journal of the Science of Food and Agriculture*, 87 (7), 1301–1307. doi: 10.1002/jsfa.2846
18. Primo-Mart n, C., Wang, M., Lichtendonk, W. J., Plijter, J. J., Hamer, R. J. (2005). An explanation for the combined effect of xylanase-glucose oxidase in dough systems. *Journal of the Science of Food and Agriculture*, 85 (7), 1186–1196. doi: 10.1002/jsfa.2107
19. Primo-Martin, C., Martinez-Anaya, M. A. (2003). Influence of Pentosanase and Oxidases on Water-extractable Pentosans during a Straight Breadmaking Process. *Journal of Food Science*, 68 (1), 31–41. doi: 10.1111/j.1365-2621.2003.tb14110.x
20. Hilhorst, R., Dunnewind, B., Orsel, R., Stegeman, P., Vliet, T., Gruppen, H., Schols, H. A. (1999). Baking Performance, Rheology, and Chemical Composition of Wheat Dough and Gluten Affected by Xylanase and Oxidative Enzymes. *Journal of Food Science*, 64 (5), 808–813. doi: 10.1111/j.1365-2621.1999.tb15917.x
21. Lysyuk, G. M., Oleynik, S. G., Zaparenko, G. V., Didenko, S. Y., Golik, O. V., Geyko, T. S. (2014). The technological aspects of emmer breed Golikovska. *Scientific Letters of Academic Society of Michal Baludansky*, 2 (5), 54–56.
22. Oleynik, S. G., Lysyuk, G. M., Zaparenko, A. V. (2015). Obosnovaniye technologicheskikh parametrov zamachivaniya zerna polby. Sovremennye technologii selskokhoziaystvennogo proizvodstva, 304–306.
23. Pogozhykh, M. I., Pak, A. O., Chekanov, M. A., Ishtvan, Ye. O., Pavliuk, I. M. (2014). Researches of system water of food raw materials by thermodynamic and molecular-kinetic methods. *Eastern-European Journal of Enterprise Technologies*, 5 (11 (71)), 42–46. doi: 10.15587/1729-4061.2014.27790
24. Drobot V. I. (2015). Tecknockimichnyi control syrovyny ta khlibobulochnykh I makaronnykh vyrobiv, Kyiv : Kondor-Vydavnytstvo, 972 p.
25. Olennikov, D. N., Tankhaeva, L. M. (2006). Metodika kolichestvennogo opredeleniya gruppovogo sostava uglevodnogo kompleksa rastitelnykh obyektor. *Khimiya rastitelnogo syrya*, 4, 29–33.
26. Matveeva, I. V., Beliavskaya, I. G. (2001). Biotechnologicheskiye osnovy prigotovleniya khleba. Moscow: DeLi print, 150.
27. Horalchuk, A. B., Pivarov, P. P., Grinchenko O. O., Pogozhykh, M. I., Polevich, V. V., Gurskyi, P. V. (2006). Rheologichni metody doslidzhennia syrovyny i kharchovykh produktiv ta avtomatyatsiya rozrakhunkiv rheologichnykh kharakterystyk. Kharkiv: KhDUHT, 63.
28. Hemdane, S., Jacobs, P. J., Dornez, E., Verspreet, J., Delcour, J. A., Courtin, C. M. (2015). Wheat (*Triticum aestivum* L.) Bran in Bread Making: A Critical Review. *Comprehensive reviews in food science and food safety*, 15 (1), 28–42. doi: 10.1111/1541-4337.12176
29. Gracheva, I. M., Krivkova, A. Yu. (2000). Technologiya fermentnykh preparatov. Moscow: «Elevar», 512.

CONSUMER PROPERTIES IMPROVEMENT OF SUGAR COOKIES WITH FILLINGS WITH NON-TRADITIONAL RAW MATERIALS WITH HIGH BIOLOGICAL VALUE (p. 54-61)

Alina Tkachenko, Inna Pakhomova

According to the study of experimental data, the feasibility of unconventional raw material with increased biological value (powdered apples, dried apricots; sea-buckthorn and calendula officinalis syrups, fat-free whey powder, whey protein concentrate, preparation of eggshells with lemon juice, cherry plums and zucchini jam, sea-buckthorn jam) and unconventional oils (pumpkin, sea-buckthorn)

in the sugar cookies production has been proved. The study indicates the increased content of essential amino acids by 34,83 % in the cookies "Dachne" and in the cookies "Yasne sonechko" by 32,54 % as compared with the check sample. The content of polyunsaturated fatty acids and minerals has increased. Especially, the content of calcium has increased by 2,8–3 times. The total fraction of carbohydrates and fats has decreased; the fraction of proteins has increased. The new cookies samples have been evaluated by the parameters of taste and smell higher than the check sample especially according to the organoleptic evaluation. The developed samples meet the standard by the physical-chemical parameters and safety parameters. The results can be used for implementation in the confectionery companies for the diversification of products with high nutritional value.

Keywords: pastry products, unconventional raw material, consumer properties, nutrition value, sugar cookies with fillings.

References

- Galushko, O. S. (2009). The trends of development market of confectionery and features of transformation in the system of values it's participants. *Actualni problemi economici*, 8, 17–25.
- Tkachenko, A. S., Syrokhman, I. V. (2015). The improvement of consumer properties of sugar cookies. *Kharchova nauka i tekhnolohiya*, 3, 82–87.
- Bodak, M. P. (2014). The usage of non-traditional raw-materials for bakery. *Visnik Lvivskoi comerciynoi academii*, 14, 113–116.
- Hadeeva, S. O., Svidlo, K. V. (2010). The determination potential risks of technology of baked biscuit semi with dietary supplements. *Zbirnik naukovih prats' NTU "KhPI"*, 46, 276–283.
- Tkachenko, A. S. (2015). The improvement of fat-acid contents of sugar cookies. *Visnik Lvivskoi comerciynoi academii*, 15, 114–119.
- Tkachenko, A. S. (2015). The sugar cookies with enriched protein contents. *Torgiylia, comerciia, marketing*, 18, 118–122.
- Shemanskaia, E. I., Oseiko, N. I. (2012). The phospholipid fat-containing products of functional appointment. *The food science*, 1, 28–30.
- Sincik, M., Goksoy, A. T. (2014). Investigation of Correlation between Traits and Path Analysis of Confectionary Sunflower Genotypes. *Notulae Botanicae Horti Agrobotanici Cluj-Napoca*, 42 (1), 227–231. doi: 10.15835/nbha4219429
- Ajay, B. C., Gowda, M. V. C., Rathnakumar, A. L., Kusuma, V. P., Fiyaz, R. A., Holajjer, P. et. al. (2011). Improving Genetic Attributes of Confectionary Traits in Peanut (*Arachis hypogaea* L.) Using Multivariate Analytical Tools. *Journal of Agricultural Science*, 4 (3), 247–258. doi: 10.5539/jas.v4n3p247
- Kulichenko, A. I. (2013). The usage of products with a milk raw material for confectionery products. *Molodoy uchenyy*, 4, 675–577.
- Nikiforova, T. A., Sevirenko, S. M., Kulikov, D. A. et. al. (2009). The prospects of usage secondary raw-materials of cereals productions. *Khleboprodukti*, 7, 50–51.
- Assel, K., Izembayeva, Meruert P. Bayisbayeva, Bayan Z., Muldabekova H. (2014). Non-Traditional Raw Materials in Production of Sugar Cookies, *American–Eurasian J. Agric. & Environ. Sci.*, 3, 358–362.
- Maximova, A. A. (2010). Innovation technology of production of the oat cookies. *Khleboprodukti*, 7, 38–39.
- Eckerson, J. M., Lieb, A. E., Mullen, K. A., Allen, N. O. (2015). The Effect of a Protein-Rich Egg White Breakfast and an Isocaloric Toaster Pastry Breakfast on Satiety and Dietary Intake in Normal-Weight College-Age Women. *Journal of the Academy of Nutrition and Dietetics*, 115 (9), A17. doi: 10.1016/j.jand.2015.06.048
- Rosliakov, I. F., Gonchar, V. V., Shulvinskaia, I. V. (2007). The usage of the Siberian pine nut kernel (*pinus sibirica*) in the production of pastry for functional appointment. *Fundamental research*, 7, 89–90.
- Rubilar, M., Gutiérrez, C., Verdugo, M., Shene, C., Sineiro, J. (2010). Flaxseed as a source of functional ingredients. *Journal of soil science and plant nutrition*, 10 (3), 373–377. doi: 10.4067/s0718-95162010000100010
- Li, S.-S., Wang, L.-S., Shu, Q.-Y., Wu, J., Chen, L.-G., Shao, S., Yin, D.-D. (2015). Fatty acid composition of developing tree peony (*Paeonia section Moutan DC.*) seeds and transcriptome analysis during seed development. *BMC Genomics*, 16 (1). doi: 10.1186/s12864-015-1429-0
- Fatima, T., Snyder, C. L., Schroeder, W. R., Cram, D., Datla, R., Wishart, D. et. al. (2012). Fatty Acid Composition of Developing Sea Buckthorn (*Hippophae rhamnoides* L.) Berry and the Transcriptome of the Mature Seed. *PLoS ONE*, 7 (4), e34099. doi: 10.1371/journal.pone.0034099
- Dorokhovich, A. M., Soloviova, O. L., Dorokhovich, V. V. (2011). Fortification of the confectionery. *Produkty & ingrediyenty*, 3, 26–28.
- Bogatirov, A. (2011). Scientific principles of fortification nutrition products. *Khlibopekarska i kondyterska promyslovist Ukrayiny*, 5, 44–47.
- Magomedov, G. O., Plotnikova, I. V., Magomedov, N. G. (2014). Beet feelings for bakery. *Khleboprodukty*, 8, 44–47.
- Aleksenko, E. V., Dikareva, I. M. (2014). The usage of see button concentrate in the cakes products. *Konditerskoe i khlebopiekarskoe proizvodstvo*, 5-6, 6–8.
- Ivanova, V. D., Khlebutina, M. S., Ivchuk, N. P. (2011). The researching of the cake functional properties. *Naukovi pratsi ONAKHT*, 40 (1), 82–86.
- Poljakova, A. V. (2011). Technology of the puff pastry with berries dry-powders supplements for puff pastry. *Visnyk Donetskoho natsionalnoho universytetu ekonomiky i torhivli imeni Mykhayla Tuhan-Baranovskoho. Ser.: Tekhnichni nauky*, 1, 55–60.
- Pop, A., Muste S., Man, S. (2013). Study of Valorification of *Lycium barbarum* (Goji) in Pastry Products / Anamaria Pop, Sevastita Muste, Simona Man. *Bulletin UASVM Food Science and Technology*, 70 (2), 93–98.
- Zucco, F., Borsuk, Y., Arntfield, S. D. (2011). Physical and nutritional evaluation of wheat cookies supplemented with pulse flours of different particle sizes. *LWT – Food Science and Technology*, 44 (10), 2070–2076. doi: 10.1016/j.lwt.2011.06.007
- Pérez, S., Matta, E., Osella, C., de la Torre, M., Sánchez, H. D. (2013). Effect of soy flour and whey protein concentrate on cookie color. *LWT – Food Science and Technology*, 50 (1), 120–125. doi: 10.1016/j.lwt.2012.06.015
- Gedrovica, P., Karklina D. (2013). Influence of Jerusalem Artichoke Powder on the Nutritional Value of Pastry Products. *Biomolecular, Agricultural, Food and Biotechnological Engineering*, 6, 7
- Aller, M. C., Villarin, M. D., Pascual, P. R. L. (2015). Product Development of Malunggay (*Moringga Oleifera*) and Sweet Potato (*Ipo-meia Batatas*) for Pastry and Fillings. *Tropical Technology Journal*, 19 (1), 5. doi: 10.7603/s40934-015-0005-1
- Recipes of the cookie (1987). Moscow: Vsesoyuznyi nauchno-issledovatel'skiy institut konditerskoy promyshlenosti, 248.
- Krutovy, Z. A., Zakharenko, G. V., Kasilova, L. O. et. al. (2013). Matematychne modeluvannya retsepturnoyi kompozitsiyi keksu pidvyshchenoyi kharchovoyi tsinnosti. *Nauka ta innovatsiyi*, 5 (9), 5–9.
- Regulation (EU) № 854 of the European Parliament and of the Council. Laying down specific rules for the organisation of official controls on products of animal origin intended for human consumption (2004). Official Journal of the European Union, 45.
- Ovchinnikova, I. A. (1974). New methods of analysis of amino acids, proteins and peptydov. Moscow: Myr, 120.

34. Otsenka nekotorykh pышчевыkh dobavok (1994). 37-y doklad Obedynennoho komyteta ekspertov FAO/VOZ po pышчевым dobavkam. Seryya tekhnicheskikh dokladov VOZ: Geneva, 27–39.
35. AACC International. Methods 10-50D and 10-52. Approved Methods of the American Association of Cereal Chemists, 10th ed. (2000). AACC International, St. Paul, MN, U.S.A.
36. Tkachenko, A. S. (2015). The forecasting of demand on the new sugar cookie with non-traditional raw-materials. Materiały IV regionalnoi konferencii: The goods of Lviv region on the today market: positioning, quality, safety within Eurointegration, 220–224.

THE EFFECT OF GRAPE SEED POWDER ON THE QUALITY OF BUTTER BISCUITS (p. 61-66)

Olga Samohvalova, Nataliya Grevtseva,
Tatiana Brykova, Anjelika Grigorenko

Butter biscuits that enjoy stable great demand in all segments of population traditionally contain a lot of fats and carbohydrates versus small amounts of biologically active substances. Enriching butter biscuits with dietary fibre, polyphenolic compounds, minerals and vitamins has a beneficial effect on the human body. These components are abundant in raw plant materials, primarily in powdered grape pomace. Grape pomace is a secondary product of wine manufacturing that is output in large quantities at wineries in Ukraine; it is an available and inexpensive raw stuff with a rich chemical composition.

We have studied the possibility of using grape seed powder in the technology of butter biscuits, in order to increase their biological value. We have found that in comparison with wheat flour, the powder is characterized by a higher water absorption capacity, and adding grape seed powder makes gluten less tensile and more elastic. The study has proved a positive effect of grape seed powder on physicochemical and organoleptic parameters of the quality of butter biscuits that become biologically more valuable.

Keywords: grape pomace, grape seed, powder, gluten, water absorption ability, butter biscuits.

References

1. Lisjuk, G. M., Vereshko, N. V., Chujko, A. M. (2011). Novi naprijamni vikoristannja vtorinnih produktiv pererobki vinogradu u virobnictvi boroshnjah virobiv. Kharkiv: HDUHT, 175.
2. Bareeva, N. N., Donchenko, L. V. Vinogradnye vyzhimki – perspektivnyj promyshlennyj istochnik pektinovyh veshhestv. Available at: <http://ej.kubagro.ru/2006/04/30/>
3. Bat'kova, I. A., Jashina, I. A., Makarova, N. V., Novikova, M. N., Smirnova, N. V. (2014). Vlijanie temperatury sushki na himicheskij sostav i antioksidantnye svojstva vinogradnyh vyzhimok. Hranenie i pererabotka sel'hozsyr'ja, 2, 36–38.
4. Holcombe, R. F., Nguyen, A. V., Martinez, M., Stamos, M. J., Moyer, M. P., Planutis, K. et. al. (2009). Results of a phase I pilot clinical trial examining the effect of plant-derived resveratrol and grape powder on Wnt pathway target gene expression in colonic mucosa and colon cancer. Cancer Management and Research, 25–37. doi: 10.2147/cmr.s4544
5. Suwannaphet, W., Meeprom, A., Yibchok-Anun, S., Adisakwattana, S. (2010). Preventive effect of grape seed extract against high-fructose diet-induced insulin resistance and oxidative stress in rats. Food and Chemical Toxicology, 48 (7), 1853–1857. doi: 10.1016/j.fct.2010.04.021
6. Salim, S., Patki, G., Jannise, B. (2015). The Role of Grape Powder in Emotional Well-Being and Memory Improvement. Diet and Nutrition in Dementia and Cognitive Decline, 925–934. doi: 10.1016/b978-0-12-407824-6.00085-9
7. Tkachenko, A. I. (2009). Vinograd – venec tvorenija prirody. Nauchno-metodicheskoe posobie dlja vrachej i shirokogo kruga chitatelej. Kharkiv: Nauchno-lechebnyj fitocentr «Avicenna», 28.
8. Sagindykova, B. A., Tihonov, A. I., Isabekova, D. S. (2011). Razrabotka tabletok s suhim jekstraktom semjan vinograda s modifitsirovannym vysvobozhdeniem. Visnik farmacii, 1 (65), 16–19.
9. Voronina, L. M., Zagajko, A. L., Samohin, A. S., Aljeksjejeva, L. M. (2004). Polifenol'ni ekstrakty vynogradu kul'turnogo na zahysti pechinky za umov oksydativnogo stresu. Klinichna farmacija, 8 (2), 36–37.
10. Grape Extracts May Be Effective Against Harmful Gut Bacteria. Available at: <https://www.sciencedaily.com/releases/2009/03/090304132621.htm>
11. Pticyn, A. R. (2007). Tehnologija vydelenija flavonoidov vinograda Vitis vinifera sorta "Izabella" dlja kosmetiki i izuchenie ih svojstv. Moscow, 26.
12. Bondakova, M. V. (2014). Razrabotka receptury i tehnologii proizvodstva kosmeticheskikh izdelij s ispol'zovaniem jekstrakta vinograda. Moscow, 23.
13. Sant'Anna, V., Christiano, F. D. P., Marczak, L. D. F., Tessaro, I. C., Thys, R. C. S. (2014). The effect of the incorporation of grape marc powder in fettuccini pasta properties. LWT – Food Science and Technology, 58 (2), 497–501. doi: 10.1016/j.lwt.2014.04.008
14. Cagdas, E., Kumcuoglu, S. (2014). Effect of grape seed powder on oxidative stability of precooked chicken nuggets during frozen storage. Journal of Food Science and Technology, 52 (5), 2918–2925. doi: 10.1007/s13197-014-1333-7
15. Aksoylu, Z., Çağındı, Ö., Köse, E. (2015). Effects of Blueberry, Grape Seed Powder and Poppy Seed Incorporation on Physicochemical and Sensory Properties of Biscuit. Journal of Food Quality, 38 (3), 164–174. doi: 10.1111/jfq.12133
16. Kalinov's'ka, T. V., Krapivnyc'ka, I. O., Obolkina, V. I., Kyjanyca, S. G. (2013). Vykorystannja vtorynnih produktiv pererobky vynogradu pid chas rozrobky innovacijnyh tehnologij kondyters'kyh vyrobiv. Obladannja ta tehnologii harchovyh vyrobnyctv. Donec'kyi nacional'nyi universitet ekonomiky i torgivli im. M. Tugan-Baranovs'kogo, 30, 75–80.
17. Obolkina, V. I., Krypichenkova, O. M., Bukshyna, L. S., Krapivnyc'ka, I. O. (2012). Pat. 72163 Ukrai'na, MPK (2012.01) A 23 G 3/00. Sposob vyrobnyctva biskvitno-zbyvnogo zdobnogo pechyva «Shantane». Zajavnyk ta patentovlasnyk Nacional'nyj universitet harchovyh tehnologij. № u201200701; zajavl. 23.01.2012; opubl. 10.08.2012, Bull. № 15, 3.
18. Zadorozhnia, O. S., Gavrysh, A. V., Docenko, V. F. (2014). Pat. 89005 Ukrai'na, MPK (2014.01) A 21 D 2/00. Pisochne pechyvo «Sonechko». Zajavnyk ta patentovlasnyk Nacional'nyj universitet harchovyh tehnologij. № u201312347; zajavl. 21.10.2013; opubl. 10.04.2014, Bull. № 7, 2.
19. Korec'ka, I. L., Lytvyn, G. V., Bandurenko, G. M., Levkiws'ka, T. M., Zinchenko, T. V. (2014). Pat. 57628 Ukrai'na, MPK (2011.01) A 23 G 3/36. Zdobne pechyvo. Zajavnyk ta patentovlasnyk Nacional'nyj universitet harchovyh tehnologij. № u201312347; zajavl. 21.10.2013; opubl. 10.04.2014, Bull. № 7, 2.
20. Jablons'ka, I. O., Myroshnyk, Ju. A., Gavrysh, A. V., Docenko, V. F. (2014). Pat. 94940 Ukrai'na, MPK (2006.01) A 21 D 13/08. Sklad pisochnogo pechyva «Aronija». Zajavnyk ta patentovlasnyk Nacional'nyj universitet harchovyh tehnologij. № u201405677; zajavl. 26.05.2014; opubl. 10.12.2014, Bjul. № 23, 2.
21. Syrohman, I. V., Turchynjak, M. K. (2008). Pat. 35281 Ukrai'na, MPK (2006) A 21 D 13/00. Zdobne pechyvo «Spokusa». Zajavnyk ta patentovlasnyk Syrohman I. V., Turchynjak M. K. № u200804704; zajavl. 11.04.2008; opubl. 10.09.2008, Bull. № 17, 3.
22. Syrohman, I. V., Lebedyneec', V. T. (2009). Asortyment i jakist' kondyters'kyh vyrobiv. Kyiv: Centr uchbovoi' literatury, 636.

23. Kalinovskaya, T. V., Obolkina, V. I. (2014). Substantiation of using wine-making secondary products as alternative raw material for confectionery industry. *Nauka i Studia*, 14 (124), 59–62.
24. Sidorenko, A. V., Vershinina, O. L., Derevenko, V. V., Shapovalova, D. V. (2011). Tehnologicheskie osobennosti prigotovlenija hlebobulochnyh izdelij obogashchennyh poroshkom iz kozhicy vinogradnyh vyzhimok. *Izvestija vuzov. Pishhevaja tehnologija*, 4, 26–28.
25. Drobot, V. I., Docenko, V. F., Arsen'eva, L. Ju., Ustinov, Ju. V., Pereguda, N. A. (1987). Issledovanie vozmozhnosti primenenija produktov iz vinograda v hlebopechenii. Kyiv, 10.
26. Davidov-Pardo, G., Moreno, M., Arozarena, I., Marín-Arroyo, M. R., Bleibaum, R. N., Bruhn, C. M. (2012). Sensory and Consumer Perception of the Addition of Grape Seed Extracts in Cookies. *Journal of Food Science*, 77 (12), 430–438. doi: 10.1111/j.1750-3841.2012.02991.x
27. Chujko, A. M., Chujko, M. M., Orlova, O. S., Jer'omenko, S. O. (2014). Research of quality of yeast-leavened dough products and shortbread using cryopowders made of herbal raw materials. *Eastren-European Journal of Enterprise Technologies*, 2 (12 (68)), 133–137. doi: 10.15587/1729-4061.2014.22416
28. Maner, S., Sharma, A. K., Banerjee, K. (2015). Wheat Flour Replacement by Wine Grape Pomace Powder Positively Affects Physical, Functional and Sensory Properties of Cookies. *Proceedings of the National Academy of Sciences, India – Section B: Biological Sciences*. doi: 10.1007/s40011-015-0570-5
29. Drobot, V. I. (2006). Laboratornyj praktykum z tehnologij hlibopekars'kogo ta makaronnogo vyrabnyctv. Kyiv: Centr navchal'noi' literatury, 341.
30. Lur'e, I. S. (1981). *Tehnologija i tehnohimicheskij kontrol' konditerskogo proizvodstva*. Moscow: Ljogkaja i pishhevaja promyshlennost', 328.